

Nearly four in 10 US HIV infections from people unaware of infection

March 18 2019



HIV infecting a human cell. Credit: NIH

Almost 40 percent of new HIV cases in the US occur because people do

not know they are infected, while a similar proportion know but are not in treatment, according to a study released Monday.

The report by the Centers for Disease Control and Prevention (CDC) is based on 2016 data and aims to bolster a strategy outlined by President Donald Trump to end the epidemic within 10 years.

The strategy has two main strands: far more widespread screening, and enabling the infected better access to treatment from the moment they test positive.

The study found that 38 percent of infections came from HIV-positive people who were unaware of their status, and 43 percent from people who knew they were infected but took no [anti-retroviral drugs](#).

The remaining infections came from people who were receiving HIV treatment but were not yet "virally suppressed."

The CDC blamed financial, social and other reasons for people not using medication, which these days typically comes in the form of a daily pill with minimal side effects.

The study said that the [infection](#) rate from the half million people in the United States who take medication and are virally suppressed—meaning they cannot pass on the disease to others—was zero.

Screening

The most at-risk group remains [homosexual men](#), with almost three-quarters of new infections coming from men having sex with men, the report said.

Five percent of infections came from intravenous drug abuse among

homosexual men, while 10 percent came from injecting drugs among the rest of the population.

Twelve percent of infections were among heterosexuals. Overall, the highest rate of transmission was among 13 to 24-year-olds.

The Trump administration has said it will invest \$291 million in the next financial year to fight HIV/AIDS, which has plateaued since 2013 to around 39,000 annual transmissions.

The goal is to reduce that number by 75 percent within five years and by 90 percent in 10 years.

Questioned about the relatively small amount of money earmarked for the multi-billion dollar task of treating HIV carriers, CDC head Robert Redfield said he was "confident that the resources that are required to accomplish this mission are in the long term plan."

The CDC, based in Atlanta, Georgia, wants doctors to make HIV screening a routine procedure.

"Everyone between the ages of 13 and 64 should get tested for HIV at least once in their lifetime," said Eugene McCray, the head of the CDC's Division of HIV/AIDS Prevention.

"Those at higher risk should get tested at least annually," he said.

"The key to controlling is helping those with HIV to control the virus," said the CDC's Jonathan Mermin, who focuses on preventing the spread of the HIV as well as other sexually transmitted diseases, tuberculosis and hepatitis.

"Time spent working closely with patients who are having trouble paying

for, picking up or taking their daily medications is time well spent"

© 2019 AFP

Citation: Nearly four in 10 US HIV infections from people unaware of infection (2019, March 18) retrieved 17 April 2024 from <https://medicalxpress.com/news/2019-03-hiv-infections-people-unaware-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.