

## Changes in onset of spring linked to more allergies across the US

March 28 2019



Credit: CC0 Public Domain

Human-induced climate change is disrupting nature's calendar, including when plants bloom and the spring season starts, and new research from the University of School of Public Health suggests we're increasingly paying the price for it in the form of seasonal allergies.



The study, based on over 300,000 respondents between 2002 and 2013, shows that seasonal allergic rhinitis, or <u>hay fever</u>, increases when the timing of <u>spring</u> "greenup" changes. The findings were published today in the journal *PLOS ONE*.

"We found that areas where the onset of spring was earlier than normal had 14% higher prevalence of hay fever," said Associate Professor Amir Sapkota in the Maryland Institute for Applied Environmental Health. "Surprisingly, we also found similar risk in areas where the onset of spring was much later than what is typical for that geographic location," he added.

The study provides the first national-level quantitative data showing how ongoing <u>climate</u> change is increasing the allergic disease burden in the United States. Hay fever, or "seasonal allergic rhinitis," affects 25 million adults in the United States and results in \$11.2 billion in related medical expenses annually.

Dr. Sapkota and his team used <u>satellite data</u> collected by NASA's Moderate Resolution Imaging Spectroradiometer (MODIS) to identify the start of the spring season throughout the US and linked these data to National Health Interview Survey data collected by the Centers for Disease Control and Prevention (CDC).

"Previous studies have shown that plant phenology is highly dependent on temperature. As such, it is one of the most sensitive indicators of how our ecosystem is responding to climate change," said Dr. Chengsheng Jiang, co-author of the paper. "We show that such climate change-driven ecological changes are directly linked to allergic disease burden in the United States. Even a relatively small change in the timing of tree flowering can have a significant economic impact given that 25 million American adults already suffer from hay fever each year."



The reason for increased hay fever when spring arrives early seems to be related to pollen exposure. An earlier onset of spring means trees flower sooner and create a longer season for tree pollen, which is the major source of spring seasonal allergens. But a very late onset of spring may mean many species of trees simultaneously burst in bloom, blasting allergy suffers with a high concentration of pollen for a shorter duration.

Dr. Sapkota suggests this is yet another example of how climate change is impacting burden of allergic disease right here in our backyard. Given that climate change is projected to intensify, Dr. Sapkota argues such data are critical to inform public health adaptation strategies, including early warning systems. "Climate change impacts our health in more ways than we can imagine. We need community-specific adaptation strategies to increase resilience and minimize disease burden associated with <u>climate change</u>," he urged.

Provided by University of Maryland

Citation: Changes in onset of spring linked to more allergies across the US (2019, March 28) retrieved 21 June 2024 from <u>https://medicalxpress.com/news/2019-03-onset-linked-allergies.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.