


Pregnancy disorder subject of new Tulane study

March 12 2019, by Barri Bronston



Dylan Lawrence, left, a graduate student in biomedical engineering, is part of assistant professor Carolyn Bayer's research team. Credit: Sally Asher

An imaging technique used to detect some forms of cancer can also help detect preeclampsia in pregnancy before it becomes a life-threatening

condition, a new Tulane study says.

Preeclampsia is a hypertensive disorder that accounts for 14 percent of global maternal deaths annually and affects 5 to 8 percent of all pregnancies. Symptoms may include [high blood pressure](#) and protein in the urine and typically occurs after the 20th week of pregnancy.

The study, published in the open-access journal *Scientific Reports* was led by Carolyn Bayer, an assistant professor of biomedical engineering, along with a team of graduate students and researchers with the Tulane School of Medicine and Tulane School of Public Health and Tropical Medicine.

The study was conducted on pregnant rats using spectral photoacoustic imaging, a noninvasive procedure that can detect placental ischemia—a sign of possible preeclampsia—prior to the onset of symptoms, such as high blood pressure, severe headaches and dizziness.

Photoacoustic [images](#) were acquired of the placenta of normal pregnant rats and rats with preeclampsia on various days of gestation. Two days after inducing preeclampsia, the average placental oxygenation decreased 12 percent in comparison to normal pregnant [rats](#).

"Spectral photoacoustic imaging is a powerful preclinical tool that has many promising applications in the understanding and treatment of pregnancy-related diseases," Bayer said.

"It provides new imaging techniques to look at the progression of the disease through gestation, which might be a better way to understand which patients need interventions to treat the [preeclampsia](#)."

Because it is a noninvasive procedure, it poses little to no risk to the fetus, compared to cordocentesis, a fetal blood sampling that is much

more dangerous.

Photoacoustic imaging may be used to detect breast, ovarian and other types of cancers.

More information: Dylan J. Lawrence et al, Spectral photoacoustic imaging to estimate in vivo placental oxygenation during preeclampsia, *Scientific Reports* (2019). [DOI: 10.1038/s41598-018-37310-2](https://doi.org/10.1038/s41598-018-37310-2)

Provided by Tulane University

Citation: Pregnancy disorder subject of new Tulane study (2019, March 12) retrieved 7 June 2024 from <https://medicalxpress.com/news/2019-03-pregnancy-disorder-subject-tulane.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.