

## Untangling the brain's life-support network

March 26 2019





Scientists at USC have created a new model of the global network within the



hypothalamus, the portion of the brain responsible for critical functions, such as thirst, hunger, and temperature regulation. They used a rat model, and they examined more than 40 years of research on the hypothalamus to develop this global network model. Credit: Joel D. Hahn, USC Dornsife College of Letters, Arts and Sciences

The portion of the brain known as the hypothalamus is small but mighty—it controls fundamental behaviors and physiology that are essential for survival. These include eating and drinking, sexual and defensive behaviors, sleep, and physiological control of things like body temperature, fluid balance, and the body's responses to stress.

A USC Dornsife-led study, published in the *Proceedings of the National Academy of Sciences* on March 25, has provided the first global <u>network</u> model—based on a highly detailed network map—of the inner workings of the <u>hypothalamus</u>. This is considered among the most critical part of the brain, as it supports life in all mammals, as well as fish, birds and many other animals.

The study is part of an ongoing effort to determine the structural organization of the mammalian <u>nervous system</u>, which the scientists refer to as <u>"The Neurome Project."</u>

The project is a pioneering effort to gain a better understanding of the brain, the most complex biological structure in the known universe. The scientists' efforts are currently focused on completing a model of the mammalian brain that is based on the connections of its gray matter regions.

Eventually, they would like to extend the model to include the connections between the entire nervous system and the body, together



called the "neurome." Such an achievement, combined with network maps for the different types of brain cells, would revolutionize research in a multitude of disciplines, from psychology to medicine.

To create the network model of the hypothalamus, the scientists conducted a rigorous analysis of connection data for each of 65 identified hypothalamic regions and spanning 40 years of brain research. Computer analysis of this data revealed a hierarchical organization comprised of subnetworks of the 65 regions of the hypothalamus on each side of the brain.

Out of 16,770 possible connections within and between the 65 regions on each side of the hypothalamus, the dataset collated by Hahn and his colleagues indicated that nearly 8,000 of them exist.

"This shows that the hypothalamus has a remarkably highly connected internal network," says Joel Hahn, assistant professor (research) of <u>biological sciences</u> at USC Dornsife College of Letters, Arts and Sciences and the study's lead author.

The computer analysis of the network also showed that two high-level subnetworks, on each side of the <u>brain</u>, are associated prominently with controlling either behavior or the body's physiology. (see graphic)

Additionally, the most highly connected regions, or hubs, were in the physiology-related control network. Hahn says this organization suggests that the hypothalamus may prioritize physiological over behavioral control.

"This organization makes sense from a survival standpoint, because behavior depends on an able body," he says.

The scientists also explored the lower-level subnetworks, which revealed



novel associations that could be relevant to several diseases, including neurodegenerative diseases such as Alzheimer's disease, and behavioral disorders that impact one or more of the functions in which the hypothalamus plays a central role.

Finally, as they developed the model, the scientists also made another discovery: Transmission of neuronal signals in the hypothalamus appears to be dominated by neuronal inhibition.

The scientists made the discovery as they analyzed genetic markers for inhibitory and excitatory (stimulatory) neurotransmission. First, they developed a map of excitation and inhibition throughout the hypothalamus. Then, they compared the map to the new network <u>model</u>.

They found that neuronal inhibition is most strongly associated with the most highly connected network nodes (hubs) of the hypothalamus. Hahn says this suggests that overall, the hypothalamus is restrained but primed for action.

He likens it to a water behind a dam released by opening or closing the floodgates.

"We don't know the functional consequences of this finding," Hahn says. "But from a mechanistic perspective, removing inhibition from a system that is primed for action achieves results fast. So, it could be beneficial for survival."

**More information:** Joel D. Hahn el al., "Macroscale intrinsic network architecture of the hypothalamus," *PNAS* (2019). <u>www.pnas.org/cgi/doi/10.1073/pnas.1819448116</u>



## Provided by University of Southern California

Citation: Untangling the brain's life-support network (2019, March 26) retrieved 13 May 2024 from <u>https://medicalxpress.com/news/2019-03-untangling-brain-life-support-network.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.