

Novel antibody may suppress HIV for up to four months

April 17 2019



Microscopic image of an HIV-infected T cell. Credit: NIAID

Regular infusions of an antibody that blocks the HIV binding site on human immune cells may have suppressed levels of HIV for up to four months in people undergoing a short-term pause in their antiretroviral

therapy (ART) regimens, according to a report published online today in *The New England Journal of Medicine*. Results of the Phase 2, open-label study indicate the antibody, known as UB-421, was safe and did not induce the production of antibody-resistant HIV. The study was supported in part by the National Institute of Allergy and Infectious Diseases (NIAID), a component of the National Institutes of Health, and United Biopharma, Inc.

The study was conducted in Taiwan and led by Chang Yi Wang, Ph.D., Chief Scientific Officer and Chairperson of United BioPharma, Inc. Twenty-nine volunteers with well-controlled HIV discontinued their normal regimens of daily oral ART at the time of their first [infusion](#) or one week later, depending on their ART regimen. Fourteen [study participants](#) received eight regular weekly infusions of UB-421, while 15 received eight higher-dose infusions every other week. At the end of the 8- or 16-week treatment period, all volunteers restarted their previous ART regimen and were evaluated in follow-up visits up to eight weeks later. Apart from a single participant who discontinued the study because of a mild skin rash, volunteers in both groups maintained HIV suppression (plasma HIV RNA levels under 20 copies/mL) throughout the treatment period in the absence of ART.

[Previous experimental infusions](#) of broadly neutralizing antibodies, or bNAbs, have suppressed HIV for about two weeks by targeting proteins on the virus itself, but the rapid mutation rate of HIV induces antibody-resistant strains that render the treatment ineffective. UB-421 theoretically avoids this possibility by blocking a stable human protein that HIV uses to infect T cells. Indeed, resistance to UB-421 was not seen in this study. Because the small study did not include a comparator group receiving a placebo infusion, further studies have been planned in Taiwan and Thailand to evaluate the safety and efficacy of UB-421 as a treatment for HIV. In [a related study](#), NIAID investigators currently are evaluating the safety of regular infusions of two highly potent bNAbs

that may prevent the development of resistant HIV strains by targeting two distinct areas of the virus.

More information: C Wang et al. Anti-CD4 Antibody UB-421 on HIV-1 Rebound after Treatment Interruption, *New England Journal of Medicine* (2019). [DOI: 10.1056/NEJMoa1802264](https://doi.org/10.1056/NEJMoa1802264)

Provided by NIH/National Institute of Allergy and Infectious Diseases

Citation: Novel antibody may suppress HIV for up to four months (2019, April 17) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-04-antibody-suppress-hiv-months.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.