

Asian nations in early tobacco epidemic: study

April 18 2019



From left, Wei Zheng, MD, PhD, Jae Jeong Yang, PhD, Danxia Yu, PhD, and colleagues are studying smoking patterns and associated deaths in Asian countries. Credit: Susan Urmy

Asian countries are in the early stages of a tobacco smoking epidemic with habits mirroring those of the United States from past decades, setting the stage for a spike in future deaths from smoking-related diseases.



That's the conclusion of researchers from Vanderbilt-Ingram Cancer Center and the Vanderbilt Epidemiology Center after analyzing 20 prospective <u>cohort</u> studies from mainland China, Japan, South Korea, Singapore, Taiwan and India.

Using long-term follow-up data from those cohorts, the study—published in *JAMA Network Open*—is the largest investigation in Asian countries of birth cohort-specific and county- or region-specific smoking patterns and their association with deaths.

Future deaths are likely to echo the pattern that occurred in the United States as the popularity of smoking increased during and after World War II, which resulted in lung cancer mortality peaking around 1990, said Wei Zheng, MD, Ph.D., Anne Potter Wilson Professor of Medicine.

"There is about a 30-year gap or incubation period for the mortality to occur," said Zheng, the study's senior author. "Smoking takes about 20 or 30 years to have this full effect on lung cancer mortality."

Tobacco control interventions may be having an effect on the smoking epidemic in some countries or areas because male smokers in the most recent birth cohort tended to quit smoking at younger ages.

"Asian countries that are richer, like Japan, South Korea and urban China, are doing a better job with this than rural China, India and other places," said Danxia Yu, Ph.D., who is co-first author of the study along with Jae Jeong Yang, Ph.D., a visiting research fellow from the Seoul National University, South Korea.

The study calls for immediate actions for all Asian countries to implement comprehensive <u>tobacco</u> control policies, including raising tobacco taxes, implementing laws for smoke-free areas, banning tobacco advertising, requiring warning labels for tobacco products and providing



help with quitting.

Older generations of Asians tended to start smoking later in life and smoke less than people in the United States did in past decades, Zheng said, but that behavior pattern is changing.

"Younger people in more recent cohorts started smoking at a younger age, and they smoked a lot more," Zheng said. "The deaths due to tobacco smoking also increased with this cohort."

The researchers classified the birth cohorts by decades, ranging from pre-1910 to 1950 or later. Smoking accounted for 12.5% of all-cause mortality in the pre-1920 birth cohort, 21.1% in the 1920s cohort and 29.3% for the cohort born in 1930 or later. Lung cancer deaths attributable to smoking, which were 56.6% among men in the pre-1920s cohort, increased to 68.4% for men born in 1930 or later.

The researchers also studied cohorts with more recent data for men and women born in later decades to analyze smoking habits.

The rate for men in mainland China who have ever smoked has increased. Among Chinese men born in 1950 or later, 79.4% of those living in urban areas had smoked, and 74.3% of those in rural areas had. Traditionally, Japanese men have had the highest rate of having ever smoked.

Women in Asia have a much lower rate for smoking. The average percentage of women smokers for all 20 cohort studies was 7.8% compared to 65.4% for men.

Asia will face a growing burden of <u>smoking</u>-related health problems unless urgent tobacco control policies are implemented, the authors concluded.



More information: Jae Jeong Yang et al, Tobacco Smoking and Mortality in Asia, *JAMA Network Open* (2019). DOI: 10.1001/jamanetworkopen.2019.1474

Provided by Vanderbilt University Medical Center

Citation: Asian nations in early tobacco epidemic: study (2019, April 18) retrieved 3 May 2024 from https://medicalxpress.com/news/2019-04-asian-nations-early-tobacco-epidemic.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.