


Brains of blind people adapt to sharpen sense of hearing, study shows

April 23 2019, by Kim Eckart



Left: Researchers began by measuring responses in the auditory cortex to find a map of how frequency responses were represented in the brain. The warm colors represent regions of the brain that showed the greatest response to low-pitched tones, while blue colors represent regions that responded more to high-pitched tones. Right: When researchers examined the range of frequencies each vertex of the brain was selective to, they found tuning tended to be narrower for blind individuals, which may underlie the enhanced ability of blind individuals to pick out and identify sounds in the environment. Credit: Kelly Chang/U. of Washington

Research has shown that people who are born blind or become blind early in life often have a more nuanced sense of hearing, especially when it comes to musical abilities and tracking moving objects in space (imagine crossing a busy road using sound alone). For decades scientists have wondered what changes in the brain might underlie these enhanced auditory abilities.

Now, a pair of research papers published the week of April 22 from the University of Washington—one in the *Journal of Neuroscience*, the other in the *Proceedings of the National Academy of Sciences*—use functional MRI to identify two differences in the brains of [blind individuals](#) that might be responsible for their abilities to make better use of auditory information.

"There's this idea that blind people are good at auditory tasks, because they have to make their way in the world without visual information. We wanted to explore how this happens in the brain," said Ione Fine, a UW professor of psychology and the senior author on both studies.

Instead of simply looking to see which parts of the brain were most active while listening, both studies examined the sensitivity of the brain to subtle differences in auditory frequency.

"We weren't measuring how rapidly neurons fire, but rather how accurately populations of neurons represent information about sound," said Kelly Chang, a graduate student in the UW Department of Psychology and lead author on the *Journal of Neuroscience* paper.

That study found that in the auditory cortex, individuals who are blind showed narrower neural "tuning" than sighted subjects in discerning small differences in sound frequency.

"This is the first study to show that blindness results in plasticity in the

auditory cortex. This is important because this is an area of the brain that receives very similar auditory information in blind and sighted individuals," Fine said. "But in blind individuals, more information needs to be extracted from sound—and this region seems to develop enhanced capacities as a result.

"This provides an elegant example of how the development of abilities within infant brains is influenced by the environment they grow up in."

The second study examined how the brains of people who are born blind or become blind early in life—referred to as "early blind" individuals—represent moving objects in space. The research team showed that an area of the brain called the hMT+—which in sighted individuals is responsible for tracking moving visual objects—shows neural responses that reflect both the motion and the frequency of auditory signals in blind individuals. This suggests that in blind people, area hMT+ is recruited to play an analogous role—tracking moving auditory objects, such as cars, or the footsteps of the people around them.

The paper in the *Journal of Neuroscience* involved two teams—one at the UW, the other at the University of Oxford in the United Kingdom. Both teams measured neural responses in study participants while participants listened to a sequence of Morse code-like tones that differed in frequency while the fMRI machine recorded brain activity. The research teams found that in the blind participants, the auditory cortex more accurately represented the frequency of each sound.

"Our study shows that the brains of blind individuals are better able to represent frequencies," Chang said. "For a sighted person, having an accurate representation of sound isn't as important because they have sight to help them recognize objects, while blind individuals only have auditory information. This gives us an idea of what changes in the brain

explain why blind people are better at picking out and identifying sounds in the environment."

The *Proceedings of the National Academy of Sciences* study examined how the brain's "recruitment" of the hMT+ region might help blind people track the motion of objects using sound. Participants once again listened to tones that differed in auditory frequency, but this time the tones sounded like they were moving. As has been found in previous studies, in blind individuals the neural responses in area hMT+ contained information about the direction of motion of the sounds, whereas in the sighted participants these sounds did not produce significant neural activity.

By using sounds that varied in frequency, the researchers could show that in blind individuals, the hMT+ region was selective for the frequency as well as the motion of sounds, supporting the idea that this region might help blind individuals track moving objects in space.

"These results suggest that early blindness results in visual areas being recruited to solve auditory tasks in a relatively sophisticated way," Fine said.

This study also included two sight-recovery subjects—individuals who had been blind from infancy until adulthood, when sight was restored via surgery in adulthood. In these individuals, area hMT+ seemed to serve a dual purpose, capable of processing both auditory and visual motion. The inclusion of people who used to be visually impaired lends additional evidence to the idea that this plasticity in the brain happens early in development, Fine said, because the results show that their brains made the shift to auditory processing as a result of their early-life blindness, yet maintains these abilities even after sight was restored in adulthood.

According to Fine, this research extends current knowledge about how

the brain develops because the team was not only looking at which regions of the [brain](#) are altered as a result of blindness, but also examining precisely what sort of changes—specifically, sensitivity to [frequency](#)—might explain how early [blind people](#) make sense of the world. As one of the [study participants](#) described it, "You see with your eyes, I see with my ears."

Both studies were funded by the National Eye Institute and the National Institutes of Health. The *Proceedings of the National Academy of Sciences* study was co-authored by Elizabeth Huber of the UW and Fang Jiang of the University of Nevada, Reno. The *Journal of Neuroscience* study was co-authored by Chang and Huber, as well as Ivan Alvarez, Aaron Hundle and Holly Bridge of the University of Oxford.

More information: Elizabeth Huber et al. Early blindness shapes cortical representations of auditory frequency within auditory cortex, *The Journal of Neuroscience* (2019). [DOI: 10.1523/JNEUROSCI.2896-18.2019](#)

Elizabeth Huber et al., "Responses in area hMT+ reflect tuning for both auditory frequency and motion after blindness early in life," *PNAS* (2019). www.pnas.org/cgi/doi/10.1073/pnas.1815376116

Provided by University of Washington

Citation: Brains of blind people adapt to sharpen sense of hearing, study shows (2019, April 23) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-04-brains-people-sharpen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.