

An EEG to assess a baby's developmental risk?

April 8 2019



Credit: CC0 Public Domain

Does exposure to stress early in life affect a baby's brain development, and is there a way to single out babies who might benefit from early intervention? A two-center study led by Boston Children's Hospital,



published today in *JAMA Pediatrics*, used brain EEGs to begin to get at these questions in an objectively measurable way. It found that infants whose mothers reported high levels of stress have a distinct pattern of brain activity as measured by EEG—at only 2 months of age.

"The EEG has been found to be exquisitely sensitive to perturbations in the environment, and thus we are not entirely surprised to see an association between stress in a mother's life and her infant's <u>brain</u> activity," says Charles Nelson, Ph.D., director of the Laboratories of Cognitive Neuroscience at Boston Children's Hospital and the study's senior investigator. "What we were surprised by, in part, was how early in life we see this association."

Tapping the EEG

EEGs, available in most pediatric practices, measure brain electrical activity, and are usually used to diagnose or evaluate people with epilepsy. But the EEG signal also contains other kinds of information about the brain. Prior studies have linked the signal's different frequency bands—alpha, beta, theta, and delta waves— with different kinds of brain activity.

"Activation of different frequency bands is associated with different aspects of cognitive functioning," explains Lara Pierce, Ph.D., a researcher in the Laboratories of Cognitive Neuroscience and the study's first author. "We think it tells us something about how the brain is developing."

The study enrolled 113 2-month-old babies in primary-care practices at Boston Children's Hospital and Children's Hospital Los Angeles serving mostly families from low-income backgrounds. The babies' mothers filled out questionnaires designed to measure their stress level. One inquired about stressful life events such as recent unemployment,



financial difficulties, housing insecurity, exposure to violence, illness, death and marital problems. The other directly asked about stress mothers were feeling.

The babies had EEGs taken while their mothers held them, and while they looked at a video showing infant toys. In all, 70 babies had usable EEG data.

Mothers' stress, babies' brains

Babies of mothers who reported feeling higher levels of stress tended to have lower power in the higher frequency EEG bands—gamma and beta—than the other infants. They also tended to have relatively higher power in the lower-frequency bands—delta and theta.

"Power in higher-frequency bands has been associated with more complex cognitive functions, such as language abilities later in childhood," Pierce explains. "We tend to see more power in lowerfrequency bands earlier in development, but elevated low-frequency power has also been observed in kids who have experienced psychosocial adversity. This has been interpreted in some cases as a sign of delayed <u>brain development</u>."

The researchers then took the opposite approach. They used a statistical technique to look for patterns in the EEG data that could indicate subgroups of children with different profiles of EEG activity. Then, they looked for environmental characteristics associated with those profiles.

That turned up a possible mitigating factor: Babies whose mothers had more years of education were significantly more likely to have an EEG profile with <u>higher power</u> across frequency bands. Pierce and her colleagues eventually want to tease out whether maternal education itself is beneficial for babies' brains, or whether other factors that enable



mothers to pursue further education are the ones that might be "protective."

Tracking brain trajectories

Prior research has found that babies start to show differences in EEGs by about 6 months of age. This is the first study showing EEG changes related to factors associated with maternal stress within a low-income context as early as 2 months, says Pierce. She and her colleagues plan to repeat the EEGs throughout the infants' first year and again at 24 and 36 months, and to see whether these patterns predict how children develop as they get a bit older.

"What we found maps pretty well to what we've seen in older kids who have experienced different types of adversity, or who live in low socioeconomic contexts," Pierce says. "We are seeing effects early in development that are associated with high levels of reported maternal stress. As we follow up, we hope that understanding what specific factors in infants' early environments impact the developing brain—and what factors might help to protect it—will ultimately help us understand how to best support families facing a variety of circumstances."

More information: JAMA Pediatrics (2019). DOI: 10.1001/jamapediatrics.2019.0492, jamanetwork.com/journals/jamap ... cle-abstract/2730067

Provided by Children's Hospital Boston

Citation: An EEG to assess a baby's developmental risk? (2019, April 8) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2019-04-eeg-baby-developmental.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.