

Fishy diagnostics for food allergy testing

April 12 2019

James Cook University scientists in Australia have found material commonly used for fish allergy testing is unreliable—potentially putting lives at risk.

A team led by Ph.D. candidate Thimo Ruethers from JCU's Australian Institute for Tropical Health and Medicine (AITHM) evaluated 26 commercially available [fish](#) preparations used for skin prick testing, the most common way of diagnosing fish allergy in Australia.

"We found the amount of allergens present varied greatly, and in some preparations major fish allergens could not be detected. This means many skin prick tests that show a person being tested is not allergic to fish may be wrong," said Mr Ruethers.

Professor Andreas Lopata from AITHM said the diagnosis of food allergy is fraught with problems.

"There are no regulations requiring the standardisation of commercial preparations for skin prick testing, while improved preparations are pricey and often impossible to get," he said.

Mr Ruethers said results from skin prick testing need to be questioned as the current preparations appear unreliable in diagnosing fish allergy.

"We urgently need reliable tools to diagnose fish allergy, a serious and often life-threatening immune disease. Current tests can be falsely negative, putting lives at risk," he said.

Professor Lopata added that seafood is now the number one cause of allergy-related deaths in Australia. He and his team are working on advanced diagnostics and therapy for seafood allergy.

Professor Lopata added that hundreds of different fish species containing different allergens are consumed in Australia, but commercial preparations for allergy testing are available for only four species.

Background:

Fish allergy is one of the most common food allergies worldwide among those that are not outgrown after childhood. Allergic reactions can be life-threatening and there is no cure for the serious, chronic immune disease. People suffering from fish allergy must avoid any food potentially containing fish.

The research team at JCU currently evaluates over 100 Australian fish species for their allergens. A recent discovery was the 'ray of hope' for fish allergy sufferers. This found that fish with a cartilaginous skeleton, including ray, are for some patients less allergenic than their fishy counterparts with bony skeletons.

Further investigations are underway, and it's hoped they will lead to improved diagnostics of fish allergy and safe fish menus for fish [allergy](#) sufferers.

More information: Thimo Ruethers et al, Variability of allergens in commercial fish extracts for skin prick testing, *Allergy* (2019). [DOI: 10.1111/all.13748](https://doi.org/10.1111/all.13748)

Provided by James Cook University

Citation: Fishy diagnostics for food allergy testing (2019, April 12) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-04-fishy-diagnostics-food-allergy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.