

Heavier and taller children are more likely to develop kidney cancer as adults than their average-sized peers

April 29 2019

A study of more than 300,000 individuals in Denmark, presented at this year's European Congress on Obesity in Glasgow, Scotland (28 April-1 May), reveals that heavier and taller children are at greater risk than their average-sized peers of developing renal cell carcinoma (RCC) as adults.

RCC is the most common form of kidney cancer found in adults. Although it often occurs in men between the ages of 50 and 70, the cancer can be diagnosed throughout adulthood. Medical experts don't know the exact causes of RCC.

"We know that overweight in adulthood is associated with an increased risk of RCC. We also know that cancers take many years to develop. We therefore had a theory that already being overweight in childhood would increase the risk of RCC later in life," explains lead author Dr. Britt Wang Jensen.

To tease out the relationships between childhood body size and the risk of RCC in adulthood, Dr. Jensen, of the Center for Clinical Research and Prevention at Bispebjerg and Frederiksberg Hospital and her colleagues used data from the Copenhagen School Health Records Register (CSHRR). The CSHRR is an electronic data base of health examination information with data from 372,636 [children](#) born in Copenhagen in the years 1930 to 1989 (and aged 30 to 89 years now). It contains serial measurements of height and weight as well as [birth weight](#)

from 1942 onwards reported by the parents.

In their study the researchers included 301,422 individuals (152,573 men) from the CSHRR, born from 1930 to 1985. The weights and heights were measured at annual school health examinations at the ages 7-13 years, and body mass index (BMI) was used to categorise the children as normal-weight or overweight, based on age- and sex- specific cut-offs suggested by the International Obesity Task Force. Cases of RCC were identified by linkage to the Danish Cancer Registry.

To analyse the data, the researchers used a statistical technique known as Cox proportional hazards regression. The procedure relates several factors or exposures—considered simultaneously—to measure the risk of an outcome, in this case, the risk of developing RCC.

During a median of 32 years of observation, 1,010 individuals (680 men) were diagnosed with RCC. Among men and women significant and positive associations were observed between childhood BMI and height, respectively, and RCC risk. When comparing two 13-year old children with one z-score difference in BMI (equivalent to 5.9 kg for boys and 6.8 kg for girls), but with similar height, the heaviest boy or girl had, in each case, a 14% higher risk of RCC than the leaner child. For height, a one z-score difference in two 13-year old children (equivalent to 8.0 cm for boys and 6.9 cm for girls) was associated with a 12% increased risk of RCC later in life for the taller boy or girl.

Compared to children with a normal-weight at 7 and 13 years, children with overweight at both ages did not have increased risks of RCC, whereas children with normal-weight at 7 years and overweight at 13 years had a 67% greater risk of developing this cancer. Compared to children with an [average height](#) at 7 and 13 years, children who were 0.5 z-score taller than average at age 7 years (equivalent to 2.6 cm for boys and girls) and remain taller than average until age 13 years have a 6%

increased risk of RCC. Children who grew from average to above average height (0.5 z-score change is equivalent to 4.0 cm additional growth in height for boys and 3.5 cm for girls) had an 8% increased risk of RCC.

The authors say: "We have found in other studies that childhood [height](#) is positively associated with several cancer forms. Therefore, we did expect to find that tall children have a higher risk of RCC than average-sized children."

They conclude: "Our findings that heavier and taller children have increased risks of RCC opens the door to new ways to explore the causes of kidney cancer."

Provided by European Association for the Study of Obesity

Citation: Heavier and taller children are more likely to develop kidney cancer as adults than their average-sized peers (2019, April 29) retrieved 2 May 2024 from <https://medicalxpress.com/news/2019-04-heavier-taller-children-kidney-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--