


Just an hour of weekly walking staves off disability

April 1 2019



Credit: CC0 Public Domain

Just one hour a week of brisk walking—as if you are late to an appointment or trying to make a train—staves off disability in older adults with arthritis pain, or aching or stiffness in a knee, hip, ankle or

foot, reports a new Northwestern Medicine study.

"This is less than 10 minutes a day for people to maintain their independence. It's very doable," said lead author Dorothy Dunlop, professor of preventive medicine at Northwestern University Feinberg School of Medicine. "This minimum threshold may motivate inactive [older adults](#) to begin their path toward a physically [active lifestyle](#) with the wide range of health benefits promoted by physical activity."

The study will be published April 1 in the *American Journal of Preventive Medicine*.

An estimated 14 million older adults in the U.S. have symptomatic knee [osteoarthritis](#), which is the most common form of osteoarthritis. Approximately two in five people with osteoarthritis—most of whom have it in their lower joints—develop disability limitations.

The study found an hour of weekly moderate-to-[vigorous physical activity](#) allowed older adults to maintain their ability to perform daily tasks like getting dressed or cross a street before a traffic light walk signal changed.

The weekly hour of exercise reduced their risk of mobility disability (walking too slowly to safely cross a street or less than one meter per second) by 85 percent and their risk of activities of daily living disability (difficulty performing morning routine tasks such as walking across a room, bathing and dressing) by almost 45 percent.

Four years after the start of the study, 24 percent of adults who did not get the weekly hour of brisk physical activity were walking too slowly to safely cross the street, and 23 percent reported problems performing their morning routine.

Study investigators analyzed four years of data from more than 1,500 adults in the national Osteoarthritis Initiative from Baltimore, Pittsburgh, Columbus and Pawtucket, Rhode Island. The adults all had pain, aching or stiffness in lower extremity joints from osteoarthritis but were free of disability when they began the study. Their physical activity was monitored using accelerometers.

"Our goal was to see what kind of activity would help people remain free of disability," Dunlop said.

Federal guidelines recommend older adults with arthritis should participate in low-impact activity. For substantial health benefits including reducing the risk for [heart disease](#) and many other chronic diseases, these guidelines recommend older adults participate in at least 2.5 hours a week of moderate-intensity activity.

But that level of activity can be daunting for inactive older adults with lower extremity pain, Dunlop said.

"We hope this new public health finding will motivate an intermediate [physical activity](#) goal," Dunlop said. "One hour a week is a stepping stone for people who are currently inactive. People can start to work toward that."

Provided by Northwestern University

Citation: Just an hour of weekly walking staves off disability (2019, April 1) retrieved 13 March 2024 from <https://medicalxpress.com/news/2019-04-hour-weekly-staves-disability.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--