

Identifying a key player in gut defense development

April 8 2019





The loss of Sox8 protein decreased the number of matured M cells (GP2, blue dots) that are responsible for establishing intestinal immune system after weaning in mice. Credit: Kimura S. et al., *Journal of Experimental Medicine*, March 15, 2019

Scientists have identified a protein critical to the immune system development and antibody production in mice, which could contribute to understanding the gut defense mechanism in infants.

A joint team from Hokkaido University and Keio University has identified a gut <u>protein</u> essential for neonatal mice to fight infections, according to research published in the *Journal of Experimental Medicine*. The insight could help understand how infants develop their own intestinal immune systems after weaning.

The gut is constantly exposed to potentially harmful bacteria that come with food and is home to naturally occurring intestinal bacteria that help fend off pathogens and produce vitamins. To maintain a healthy balance, microfold <u>cells</u> or 'M' cells in the gut lining absorb antigens, which are foreign molecules from outside the body. This absorption triggers production of <u>antibodies</u>, which bind to and neutralize pathogens.

Infants do not have their own immune systems and receive maternal antibodies through breastfeeding to keep harmful bacteria at bay. However, it has been unclear how <u>young children</u> develop M cells and antibodies during the weaning phase.

In the current study, a group led by Shunsuke Kimura of Hokkaido University, and Nobuhide Kobayashi and Koji Hase of Keio University found that a protein called Sox8 was specifically expressed by M cells. Using florescence and imaging techniques, they also found that the



protein was present in both young and mature M cells, indicating its consistent involvement during the cell development stages.

Furthermore, the team found that Sox8 directly promotes the production of an antigen receptor, GP2, that is only found in fully matured M cells and enables antigen uptake.

Conversely, lack of Sox8 in mice led to reduced expression of GP2 and therefore reduced antigen uptake, suggesting that Sox8 is responsible for the maturation of M cells. In the mice lacking Sox8, production of antibodies was significantly reduced one week after weaning, indicating that the protein is indispensable for the immediate establishment of immune system after weaning.

"Future studies should investigate genes regulated by this protein in order to completely understand the molecular mechanisms underlining the intestinal immune system," says Shunsuke Kimura. "We hope that our study may one day help develop <u>preventive measures</u> such as vaccines against gut infections among infants," Koji Hase added.

More information: Shunsuke Kimura et al. Sox8 is essential for M cell maturation to accelerate IgA response at the early stage after weaning in mice, *The Journal of Experimental Medicine* (2019). DOI: 10.1084/jem.20181604

Provided by Hokkaido University

Citation: Identifying a key player in gut defense development (2019, April 8) retrieved 17 July 2024 from <u>https://medicalxpress.com/news/2019-04-key-player-gut-defense.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.