

Researchers discover why men are more likely to develop liver cancer

April 3 2019



Raising adiponectin levels (right) reduces the formation of liver tumors in male mice. Credit: Manieri et al., 2019

Researchers in Spain have discovered that a hormone secreted by fat cells that is present at higher levels in women can stop liver cells from becoming cancerous. The study, which will be published April 3 in the *Journal of Experimental Medicine*, helps explain why hepatocellular carcinoma (HCC) is more common in men, and could lead to new treatments for the disease, which is the fourth leading cause of cancer-related death worldwide.

In the United States, men are twice as likely as women to develop HCC.



A potential contributor to this <u>gender disparity</u> is adiponectin, a hormone secreted by <u>fat cells</u> that helps control the body's metabolism. "Circulating adiponectin levels have been reported to be higher in women than in men," explains Dr. Guadalupe Sabio of the Spanish National Center for Cardiovascular Research (CNIC) in Madrid. "However, adiponectin's role in HCC is controversial and needed further investigation."

Similar to humans, male mice are more prone to HCC than females. Sabio and colleagues found that increased levels of adiponectin in female mice protect them from HCC. The researchers determined that the hormone activates two proteins inside <u>liver cells</u>, known as p38 and AMPK, that block cell proliferation and impair <u>tumor growth</u>.

Inhibiting testosterone production in male rodents increased their adiponectin levels and reduced tumor growth. Sabio and colleagues found that testosterone activates a protein in fat cells called JNK1 that inhibits adiponectin production.







Raising adiponectin levels (right) reduces the formation of liver tumors in male mice. Credit: Manieri et al., 2019

"Our results unravel a clear crosstalk between sex hormones, <u>adipose</u> <u>tissue</u>, and the liver, clarifying the mechanism underlying gender disparity in liver cancer development," Sabio says.

Intriguingly, adiponectin levels are also reduced in obesity, another major risk factor for HCC. The study also suggests that adiponectin and metformin—a common antidiabetic drug that activates AMPK—could be used as novel treatments for liver cancer.

More information: Manieri et al., 2019. *J. Exp. Med.* DOI: <u>10.1084/jem.20181288</u>

Provided by Rockefeller University Press

Citation: Researchers discover why men are more likely to develop liver cancer (2019, April 3) retrieved 8 May 2024 from <u>https://medicalxpress.com/news/2019-04-men-liver-cancer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.