

## New research identifies microbes that may reduce allergy-like reactions in many people

April 16 2019



Cheese rind is removed from a wheel of ripening cheese. Iowa State University scientists are studying the bacteria in cheese rinds in an effort to reduce unpleasant reactions in people with histamine susceptibility. Credit: Monika Dzieciol

Some aged cheeses cause allergy-like reactions in many people, but an Iowa State University scientist is working to identify bacteria that could reduce those unpleasant side effects.

The answer to alleviating the unpleasant reactions that result from eating ripened cheeses could lie in the <u>bacteria</u> that populate the cheese's rind,



said Stephan Schmitz-Esser, associate professor of animal science. Schmitz-Esser led a new study, published Tuesday in the peer-reviewed <u>academic journal</u> *Scientific Reports*, that looks closely at the microbial communities that develop in cheese rinds, and the research could pave the way to improved cheese production techniques that cut down on the compound that causes those reactions.

Many hard cheese varieties, such as Comte and Gruyere, require months to ripen to obtain the desired flavor and aroma. As the cheeses ripen, a biofilm of bacteria and fungi develops on the surface of the cheese. These microbial communities, which form naturally and are not inoculated by cheese producers, play an essential role in the ripening process and can protect the cheese from harmful pathogens, Schmitz-Esser said.

"We still know surprisingly little about these <u>microbial communities</u> that grow on the cheese rind," he said. "We know that they work, but we don't know a lot beyond that."

These microbes produce histamine, a compound involved in local immune responses, as a byproduct of fermentation, he said. Foods that contain high levels of histamine can cause rashes and other symptoms associated with <u>allergic reactions</u> in roughly 1% of humans.

So Schmitz-Esser and his collaborators set out to answer some of the questions cheese producers have about those microbes. The researchers sampled cheese rinds originating in Austria and isolated the bacteria strains they contain. They identified a particular bacterial genus, called Brevibacterium, and sequenced the genomes of three different species. That data allowed them to identify pathways in the bacterial genomes that are likely to govern the degradation of histamine. Finally, the researchers grew the bacteria in an experimental medium and demonstrated their ability to break down histamine.



Schmitz-Esser said not all the Brevibacterium strains the team studied possess the genetic pathway that most likely reduces histamine content, but it may be possible to harness the strains that do express the pathway in the ripening process.

"Ultimately, this could allow us to intentionally use such strains that have the ability break down histamine," he said. "These bacteria would support the ripening process and, at the same time, reduce histamine production."

The research could allow diners who are susceptible to <u>histamine</u> reactions to enjoy ripened cheese, and it also could help cheese producers standardize the ripening process and improve efficiency, Schmitz-Esser said. The time it takes for the <u>cheese</u> to mature represents a major investment by producers. Optimizing the process ensures producers get the best return on that investment, he said.

Provided by Iowa State University

Citation: New research identifies microbes that may reduce allergy-like reactions in many people (2019, April 16) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2019-04-microbes-allergy-like-reactions-people.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.