


The protein p38-gamma identified as a new therapeutic target in liver cancer

April 10 2019



Hepatocytes during the cell division process. Credit: CNIC

An enzyme involved in cell stress responses could become a new



pharmacological target in the treatment of liver cancer, a disease that currently has few available treatment options. The research team at the Centro Nacional de Investigaciones Cardiovasculares (CNIC) led by Guadalupe Sabio has discovered that the protein p38γ, one of the four types of p38 kinase, is essential for the initiation of cell division in liver cells. This indicates that "p38γ could be a useful therapeutic target for liver cancer," says Sabio, adding, "We are now developing inhibitors of this protein to test in this cancer." The study has been published today in *Nature*.

National data on cancer rates in Spain for 2017 show that <u>liver cancer</u> is detected in an estimated 5,862 people every year (4252 men and 1610 women), making it the seventh-most common cancer among men and the 17th most common among women. In line with figures for the European Union, the incidence of liver cancer in Spain is approximately 12 in every 100,000 men and 3.5 in every 100,000 women.

Sabio has a long-standing interest in the p38 family of proteins, which are activated in <u>cells</u> exposed to stress. Now, in the study published in *Nature*, her team has shown an important role for p38 γ in the initiation of <u>cell division</u> in the liver.

"The four members of the p38 kinase family are so similar that at first they appear to have overlapping or redundant functions," says Sabio. "But detailed analysis of their three-dimensional structures thanks to collaborations with Silvia Osuna at the University of Girona and investigators at the Centro Nacional de Investigaciones Oncológicas (CNIO) revealed that one of the four, p38γ, also shares close similarities with another family of proteins called CDKs. These proteins are well-known regulators of cell division and the cell cycle and play a well-established role in the development of cancer."

In laboratory studies exploring this similarity to CDKs, the team found



that an inhibitor of CDK2 also reduced the activity of p38 γ . The researchers also found that p38 γ and CDK2 both act on a tumor suppressor protein that plays a central role in the regulation of the cell cycle.

To test whether p38γ is implicated in cell division, study first author Antonia Tomás-Loba examined the outcome of chemically inducing liver cancer in mice that lack the enzyme. The results were truly promising: "in mice lacking p38γ or treated with inhibitors to block its activity, the development of hepatocellular carcinoma was slowed," says Tomás-Loba.

These results, says Sabio, "could be extrapolated to human patients." Indeed, work with colleagues at Salamanca University Hospital shows that the amount of p38γ increases with liver fibrosis, a process that precedes cancer and is much higher in liver cancer patients. These results suggest that in the future it may be possible to treat this type of cancer with drugs that specifically target p38γ. The advantage of targeting p38γ is that this enzyme appears to control the initiation of the cell cycle in response to stress, and therefore inhibiting this process would not affect tissues that are constantly proliferating, like the intestinal lining or hair follicles.

More information: Antonia Tomás Loba et al. p38γ is essential for cell cycle progression and liver tumorigenesis, *Nature* . 10.1038/s41586-019-1112-8

Provided by Centro Nacional de Investigaciones Cardiovasculares Carlos III (F.S.P.)

Citation: The protein p38-gamma identified as a new therapeutic target in liver cancer (2019,



April 10) retrieved 25 April 2024 from https://medicalxpress.com/news/2019-04-protein-p38-gamma-therapeutic-liver-cancer.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.