

Reduction in autism diagnoses observed with DSM-5

April 2 2019



(HealthDay)—Use of the Diagnostic and Statistical Manual 5 (DSM-5)

criteria for autism spectrum disorder (ASD) seems to have reduced the number of ASD diagnoses, according to a review published online March 9 in the *Journal of Autism and Developmental Disorders*.

Kristine M. Kulage, M.P.H., from the Columbia University School of Nursing in New York City, and colleagues examined changes in the frequency of ASD diagnoses since the DSM-5 publication in a five-year follow-up systematic review and meta-analysis. Data were included for 33 studies.

The researchers found that use of DSM-5 criteria indicates reductions in the diagnosis of ASD (20.8 percent; P

"Our findings provide further insight regarding how the DSM-5 is being used nationally and internationally to diagnose, or failing to diagnose, those with ASD," Kulage said in a statement. "Future research is needed, as concerns remain for impaired individuals, who, because of the change in diagnostic criteria for ASD, may no longer qualify for treatment but still demonstrate a need for services."

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Reduction in autism diagnoses observed with DSM-5 (2019, April 2) retrieved 5 May 2024 from <https://medicalxpress.com/news/2019-04-reduction-autism-dsm-.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--