

Urine test could prevent cervical cancer

April 29 2019, by Mike Addelman



Credit: CC0 Public Domain

Urine testing may be as effective as the smear test at preventing cervical cancer, according to new research by University of Manchester scientists.

The study, led by Dr. Emma Crosbie and published in *BMJ Open*, found

that urine testing was just as good as the [cervical smear](#) at picking up high-risk human papillomavirus (HPV), the virus that causes cervical [cancer](#).

The research team say a urine [test](#) could help increase the numbers of [women](#) who are screened for cervical cancer, which affects more than 3,000 women every year in the UK.

The research was supported by the Manchester NIHR Biomedical Research Centre.

Urine testing could also have a role in the [developing world](#), where cervical cancer is up to 15 times more common and smear testing largely non-existent.

The NHS cervical screening programme tests for so called 'high-risk' types of human papillomavirus (HPV) and the health of the cells of the cervix in women who test high-risk HPV positive.

Around one in 20 women show abnormal changes which might go on to become cancer and are referred for colposcopy, where the cervix is examined under magnification, allowing abnormal areas to be seen, sampled and treated, before they ever cause cancer.

According to the team, cervical smear samples, self-collected vaginal samples and [urine samples](#) are all effective at picking up high risk HPV infection.

Cervical cancer is most common in women aged 30 to 35 years. But the precancerous stage is detectable in the five to 10 years before this, when up to a third of women fail to attend for their [smear test](#).

"We're really very excited by this study, which we think has the potential

to significantly increase participation rates for cervical cancer screening in a key demographic group," said Dr. Emma Crosbie.

"Many [younger women](#) avoid the NHS cervical cancer screening programme because they find it embarrassing or uncomfortable, particularly if they have gynaecological conditions like endometriosis."

She added: "Campaigns to encourage women to attend cervical screening have helped. The brilliant campaign by the late Jade Goody increased numbers attendance by around 400,000 women.

"But sadly, the effects aren't long lasting and participation rates tend to fall back after a while. We clearly need a more sustainable solution."

Of the 100 or so types of HPV, some are linked to cervical cancer, and some are linked to other conditions, like genital warts.

Most cervical cancers are caused by high-risk types HPV-16 and HPV-18.

104 women attending the colposcopy clinic at St Mary's Hospital, Manchester participated in the study and were screened using two brands of HPV testing kits.

Around two-thirds of the women tested positive for any high-risk HPV type, and a third for HPV-16 or HPV-18.

From the total, 18 women had pre-cancerous changes to the cervix that needed treatment.

With the Roche HPV testing kit, urine, vaginal self samples and cervical smears picked up 15 of these.

With the Abbott HPV testing kit, urine picked up 15 of these and vaginal self samples and cervical smears picked up 16.

Dr. Crosbie said: "These results provide exciting proof of principle that [urine](#) HPV testing can pick up cervical pre-cancer cells, but we need to trial it on a greater number of women before it can be used in the NHS. We hope that is going to happen soon.

"Urine is very simple to collect and most hospitals in the developed and developing world have access to the lab equipment to process and test the samples.

"Let us hope this is a new chapter in our fight against [cervical cancer](#), a devastating and pernicious disease."

Provided by University of Manchester

Citation: Urine test could prevent cervical cancer (2019, April 29) retrieved 13 May 2024 from <https://medicalxpress.com/news/2019-04-urine-cervical-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--