

Women underreport prevalence and intensity of their own snoring

April 23 2019



A new study of adults who were referred for evaluation of a suspected sleep disorder suggests that women tend to underreport snoring and underestimate its loudness. Credit: American Academy of Sleep Medicine

A new study of adults who were referred for evaluation of a suspected sleep disorder suggests that women tend to underreport snoring and underestimate its loudness.

Results show that objectively measured snoring was found in 88% of the [women](#) (591 of 675), but only 72% reported that they snore (496 of 675). In contrast, objective snoring (92.6%) and self-reported snoring (93.1%) were nearly identical in men. The study also found that women

snored as loudly as men, with a mean maximal snoring intensity of 50 decibels among women and 51.7 decibels among men. About 49% of the women had severe or very severe snoring (329 of 675), but only 40% of the women rated their snoring at this level of severity (269 of 675).

"We found that although no difference in snoring intensity was found between genders, women tend to underreport the fact that they snore and to underestimate the loudness of their snoring," said Nimrod Maimon, MD, MHA, principal investigator and professor at the BGU Faculty of Health Sciences and the head of internal medicine (B Ward) at Soroka University Medical Center in Be'er Sheva, Israel. "Women reported snoring less often and described it as milder."

The study results are published in the March 15 issue of the *Journal of Clinical Sleep Medicine*.

Snoring is a respiratory sound generated in the upper airway during sleep. The intensity of snoring may vary and often will disturb the bed partner's sleep. Snoring is a common warning sign for obstructive sleep apnea, a chronic disease that involves the repeated collapse of the upper airway during sleep.

The study involved 1,913 patients who were referred to a sleep disorders center at a university hospital for a sleep evaluation. They had an average age of 49 years. Participants were given a questionnaire that asked them to rate the severity of their snoring. Objective snoring volume was quantified using a calibrated digital sound survey meter during a sleep study that lasted an entire night. Snoring [intensity](#) was classified as mild (40—45 decibels), moderate (45—55 decibels), severe (55—60 decibels), or very severe (60 decibels or more).

According to the authors, there is a [social stigma](#) associated with snoring among women. Therefore, women may not reliably answer questions

about snoring, which may contribute to the underdiagnosis of obstructive sleep apnea in women.

"The fact that women reported snoring less often and described it as milder may be one of the barriers preventing women from reaching sleep clinics for a sleep study," Maimon said.

Maimon added that [health care providers](#) who are screening women for suspected [obstructive sleep apnea](#) should consider other factors in addition to self-reported snoring. For example, women with sleep apnea may be more likely than men to report other symptoms such as daytime fatigue or tiredness.

More information: Roi Westreich et al, The Presence of Snoring as Well as its Intensity Is Underreported by Women, *Journal of Clinical Sleep Medicine* (2019). [DOI: 10.5664/jcsm.7678](https://doi.org/10.5664/jcsm.7678)

Provided by American Academy of Sleep Medicine

Citation: Women underreport prevalence and intensity of their own snoring (2019, April 23) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2019-04-women-underreport-prevalence-intensity.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--