

Blood pressure and glucose control may prevent common arrhythmia

May 24 2019



Credit: CC0 Public Domain

Blood-pressure and glucose control may be effective in preventing heart block, a common form of arrhythmia, and the subsequent need for a pacemaker, according to a study by researchers at UC San Francisco.

In an analysis of more than 6,000 Finnish patients, appearing online May 24, 2019, in *JAMA Network Open*, UCSF researchers found that more

than half of the cases of heart block resulted from [high blood pressure](#) or elevated [blood sugar](#).

Atrioventricular (AV) block occurs when electrical conduction is impaired between the heart's four chambers, most often by fibrosis or sclerosis. It is often felt as the heart skipping a beat.

An estimated 3 million people worldwide have pacemakers, and 600,000 pacemakers are implanted annually. But while a common treatment and low-risk procedure, it can result in serious complications. Generator charges also carry a high risk of infection in and around the heart. However, there has been limited research on whether behavioral modifications can prevent heart block and which ethnicities are most at risk.

"It is perhaps precisely because pacemakers so successfully and immediately address cases of heart block that we have previously failed to devote more attention to prevention of this important disease," said senior author Gregory Marcus, MD, MAS, a UCSF Health cardiologist and associate chief of cardiology for research in the UCSF Division of Cardiology. "In addition to the prevention and treatment of myocardial infarction and heart failure, effective treatment of hypertension and maintenance of normal blood sugars may be useful prevention strategies."

This is the first community-based study to evaluate the possible association between common modifiable cardiovascular risks and heart block occurrence requiring pacemaker implantation.

In the *JAMA Network Open* study, Marcus and his colleagues used the Mini-Finland Health Survey, which was designed to represent the country's population aged 30 and over, and consisted of an in-home interview and clinic examination on various health subjects. They

studied 6,146 Caucasian patients enrolled from 1978 to 1980, then reviewed the patients' hospital records from 1987 to 2011 to determine heart block incidents.

Over an average follow up of 25 years, 58 patients developed AV block. The researchers found that older age, being male, higher [systolic blood pressure](#), higher fasting glucose, history of myocardial infarction and history of congestive [heart failure](#) independently increased the likelihood of occurrence.

Of those factors, two directly modifiable risk factors were identified: every 10 millimeter increase in systolic blood pressure resulted in 22 percent greater risk, and every millimeter increase in fasting glucose resulted in 19 percent greater risk. Taking into account the prevalence of these modifiable risk factors in the population and assuming causal relationships, they estimated that 47 percent of AV blocks in the 58 patients would have been avoided with ideal blood pressure and 11 percent with normal fasting glucose.

The authors note that the study occurred in a solely Caucasian population and advised caution in applying findings to other populations.

"Given the prevalence of [heart](#) block in the adult male population, as well as the multiple risks associated with pacemakers, it would be worthwhile to pursue further research on this connection," said Marcus, who holds the Endowed Professorship of Atrial Fibrillation Research in the UCSF School of Medicine. "This new information also may help persuade hypertensive individuals to receive and continue their prescribed treatments."

Provided by University of California, San Francisco

Citation: Blood pressure and glucose control may prevent common arrhythmia (2019, May 24)
retrieved 26 April 2024 from
<https://medicalxpress.com/news/2019-05-blood-pressure-glucose-common-arrhythmia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.