

## **Researchers find new brain mechanisms regulating body weight**

## May 3 2019

Researchers at the University of Gothenburg, Sweden, have clarified the link between the molecule interleukin-6 (IL-6) in the brain and obesity. In experiments on rats and mice, they show that the molecule does affect the risk of obesity, and also where this effect occurs in the brain.

interleukin-6 (IL-6) is a well-known pro-inflammatory molecule, and an integral element of body's first line of defense during infection. Intriguingly, the brain may govern and utilize IL-6 differently from the rest of the body. Researchers led by the laboratory of Karolina Skibicka at the Sahlgrenska Academy, Sweden wondered what happens to IL-6 levels in the brain following a diet that leads to obesity.

Rats and mice were offered a high-calorie palatable food, a mix of fat and sugar, in addition to their regular low-calorie diet. Like many humans, rodents choose to overeat when presented with calorie-dense foods.

"What we found was that the rats and mice that became obese had reduced IL-6, but only in one brain region called the lateral parabrachial nucleus (IPBN)," says Devesh Mishra, postdoctoral fellow leading the study. "To understand whether this reduction of IL-6 is a good or bad thing for the metabolic health of the rodents, we viro-genetically reduced IL-6 levels very selectively in the IPBN; this led to increased body weight and body fat, even in rodents fed a healthy diet."

Therefore, the researchers have concluded that the reduced levels of



IPBN IL-6 in obesity are problematic, and likely contribute to metabolic dysfunction and weight gain. Since <u>body weight</u> is a result of the amount of food consumed, i.e., <u>energy intake</u>, versus how much <u>energy</u> we use, i.e. <u>energy expenditure</u>, weight gain can follow dysfunction of either one of these branches of energy balance.

What makes local parabrachial nucleus-produced IL-6 extra important, the study found, is that it affects both branches simultaneously: It decreases food intake and increases energy expenditure, the latter by increasing brown fat activity, so body's energy is utilized for heat generation or fat burning. Hence, reduced levels of IPBN IL-6 disrupt the entire energy balance equation.

These findings may be relevant not only for mice but also men, since an <u>earlier study</u> from University of Gothenburg revealed that blood serum IL-6 levels in obese and overweight men are increased, yet brain IL-6 levels measured in cerebrospinal fluid are reduced.

There is one unresolved issue related to these findings—researchers found that the obesity-associated reduction in IL-6 was only present in males. Female rats and mice had normal IL-6 levels. The Sahlgrenska team is now investigating why females are protected from the obesityassociated IL-6-driven dysfunction.

Given that obesity is a major global disorder with 1.9 billion overweight individuals out of which 650 million are obese, effective anti-obesity treatments are desperately needed to minimize the amount of personal and medical burden on individuals and societies. On the scientific end, researchers think that IL-6 as a satiety-mediating substance with <u>brain</u> region specificity is an important discovery, and can open new directions in the quest for more effective anti-obesity strategies.

More information: Devesh Mishra et al. Parabrachial Interleukin-6



Reduces Body Weight and Food Intake and Increases Thermogenesis to Regulate Energy Metabolism, *Cell Reports* (2019). <u>DOI:</u> <u>10.1016/j.celrep.2019.02.044</u>

## Provided by University of Gothenburg

Citation: Researchers find new brain mechanisms regulating body weight (2019, May 3) retrieved 26 April 2024 from <u>https://medicalxpress.com/news/2019-05-brain-mechanisms-body-weight.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.