

CDC: Racial disparity seen with lupusrelated deaths

May 15 2019



(HealthDay)—Mortality from systemic lupus erythematosus (SLE) is significantly higher among blacks, according to research published in the May 10 issue of the U.S. Centers for Disease Control and Prevention *Morbidity and Mortality Weekly Report*.



S. Sam Lim, M.D., from Emory University in Atlanta, and colleagues analyzed data from the population-based Georgia Lupus Registry (2002 to 2004). Incident and prevalent SLE cases were matched to the National Death Index through 2016.

The researchers identified 97 incident case deaths and 401 prevalent case deaths. Adjusting for age group, sex, and race, standardized mortality ratios were two to three times higher among persons with SLE compared with expected deaths in the general population. There was significantly higher cumulative mortality among blacks than whites. For both incident and prevalent cases, blacks were significantly younger at death (mean age, 51.8 and 52.3 years, respectively) than whites (mean age, 64.4 and 65.0 years, respectively). Mortality among incident cases in whites did not occur until five years after SLE diagnosis, while blacks had significantly and persistently higher mortality rates from the time of diagnosis. By sex, there were no significant differences.

"The effect of this racial disparity in mortality is further underscored by the fact that the prevalence of SLE in blacks is three times that in whites," the authors write.

More information: <u>Abstract/Full Text</u>

Copyright © 2019 HealthDay. All rights reserved.

Citation: CDC: Racial disparity seen with lupus-related deaths (2019, May 15) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2019-05-cdc-racial-disparity-lupus-related-deaths.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.