

Combination therapy advisable for IBS

May 15 2019, by Elin Lindström Claessen



Magnus Simren, Professor at Sahlgrenska Academy, University of Gothenburg.
Credit: Cecilia Hedstrom

The more abnormalities in intestinal and brain function that IBS sufferers have, the more severe their symptoms of this functional bowel disorder, and the more adversely their everyday life is affected. This is shown by a Sahlgrenska Academy study indicating that patients with IBS should get treatments for different abnormalities simultaneously, to

improve both bowel function and signaling from the brain to the gut.

IBS (irritable bowel syndrome) is very common. Up to one out of ten adults in Sweden has this functional bowel disorder to some degree of severity. Diagnostic criteria for the IBS diagnosis include [abdominal pain](#), constipation or diarrhea, and irregular bowel activity over a long period. The underlying causes are not entirely known, but abnormalities both locally in the intestines and in the central nervous system are thought to be implicated.

Currently, no available treatment cures IBS, but its symptoms can be alleviated. There are drugs that can improve [intestinal function](#) in various ways, and in some cases [brain](#)-directed therapy is also given.

"There are studies showing that hypnosis, [cognitive behavioral therapy](#) (CBT) and antidepressants can all have an effect against IBS. The view of the gut and brain working together in IBS is beginning to be increasingly accepted, so many gastroenterology clinics are striving to work holistically, in multiprofessional teams, to manage their IBS patients," says Professor Magnus Simren of Sahlgrenska Academy, University of Gothenburg, Sweden, who led the work on the new study.

The study, published in the journal *Gastroenterology*, comprised 400 people with IBS and also healthy controls. The scientists measured intestinal abnormalities relating to sensitivity and motility, and the [study participants](#) were asked to reply to a questionnaire to capture signs of abnormal functioning of the central nervous system.

"We investigated several abnormalities at various levels in the nerves that connect the intestines and the brain, which is known as the 'gut-brain axis.' The questionnaires used are established for demonstrating incidence of anxiety and depression, which in our study served as a marker for abnormal functioning in the central nervous system," says

Magnus Simren.

The study shows that the people who had many different abnormalities considered to have a bearing on the symptoms of IBS, in both the [gastrointestinal tract](#) and the brain, were also those patients who reported the most severe symptoms and experienced the lowest quality of life. The association was linear: as the number of abnormalities rose, a gradual worsening in symptom severity was also observed.

"In the study, it's striking how the burden of disease increases successively the more abnormalities the patient has. This means that we must probably focus the treatment on several of these at the same time to get a better effect. So this implies that a combination of different treatments tackling factors in the gut and the brain can have a positive effect. But this hypothesis must be investigated in clinical trials first," Simren says.

More information: Magnus Simrén et al, Cumulative Effects of Psychologic Distress, Visceral Hypersensitivity, and Abnormal Transit on Patient-reported Outcomes in Irritable Bowel Syndrome, *Gastroenterology* (2019). [DOI: 10.1053/j.gastro.2019.04.019](https://doi.org/10.1053/j.gastro.2019.04.019)

Provided by University of Gothenburg

Citation: Combination therapy advisable for IBS (2019, May 15) retrieved 27 April 2024 from <https://medicalxpress.com/news/2019-05-combination-therapy-ibs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.