


A new way to predict complications after larynx cancer surgery

May 31 2019, by Nicole Fawcett



Rebecca Hoesli, M.D., and Matthew Spector, M.D., evaluate an image from the study Credit: University of Michigan Rogel Cancer Center

A technique that illuminates blood flow during surgery predicted which head and neck cancer patients were likely to have issues with wound


healing. It could enable surgeons to make adjustments during surgery or recovery to improve outcomes.

A team of surgeons at the University of Michigan Rogel Cancer Center found the approach so successful in a clinical trial that they closed the study early.

Most people with larynx cancer will have radiation and chemotherapy. But about one-third of the time, the cancer will return or will prove resistant, leaving surgery as the next option.

At this point, [tissue damage](#) from the radiation adds challenges to the operation. When the surgeon closes the wound, damaged tissue can interfere. For about 40% of patients, this will lead to a pharyngocutaneous fistula, a hole in the neck where saliva can leak out. It can cause bleeding or infections, keeping patients in the hospital longer, and in 10% of cases sending them back to the operating room to fix it.

"Radiation damage is something you can't always see. There have been very few examples in the literature that would explain or predict who's going to have a complication," says Matthew E. Spector, M.D., assistant professor of otolaryngology-head and neck surgery at Michigan Medicine. Spector is the senior author on a paper made available online in February ahead of final print publication in May in *Annals of Surgical Oncology*.



Visual abstract describes the paper's findings. Credit: University of Michigan Rogel Cancer Center

Researchers enrolled 41 patients who were undergoing laryngectomy after radiation. After removing the tumor but before closing the throat, anesthesiologists gave the patients an intravenous injection of a type of medical dye, indocyanine green. The dye circulates within about 40 seconds. Surgeons then use laser angiography, which illuminates the dye, allowing them to observe [blood flow](#).

The results were clear-cut: patients with lower blood flow had a significantly higher risk of developing a fistula, whereas patients with high blood flow had a very low risk of wound complications.

Knowing this, Spector suggests a few possible interventions. One could be cutting out a wider margin of tissue to get a cleaner, healthier edge.

Another possibility is to keep high-risk patients in the hospital longer, while sending the low-risk patients home more quickly.

The laser angiography approach would be straightforward to implement in many settings. It's already used by other [surgeons](#), including in breast reconstruction, so many hospitals already own the equipment. The technique has little impact on [patients](#) because it can be administered so quickly while they are still under anesthesia. Reactions to the indocyanine are minimal.

Researchers are developing a randomized clinical trial to assess whether cutting back more tissue leads to fewer fistulas in the high-risk group.

"We need to find an intervention that can lower this risk," Spector says.

More information: Rebecca Hoesli et al, Intraoperative Fluorescent Angiography Predicts Pharyngocutaneous Fistula After Salvage Laryngectomy, *Annals of Surgical Oncology* (2019). [DOI: 10.1245/s10434-019-07262-6](#)

Provided by University of Michigan

Citation: A new way to predict complications after larynx cancer surgery (2019, May 31) retrieved 18 April 2024 from <https://medicalxpress.com/news/2019-05-complications-larynx-cancer-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--