


How the dengue virus replicates in infected cells

May 9 2019

Amino acid residues in NS1 required for:



Overview of amino acid residues in NS1 required for NS1 secretion, interaction with the NS4A-2K-4B precursor and RNA replication. Credit: Plaszczyca A, *et al.* (2019)

The nonstructural protein 1 (NS1) of the dengue virus interacts with another viral protein called NS4A-2K-4B to enable viral replication, according to a study published May 9 in the open-access journal *PLOS Pathogens* by Ralf Bartenschlager of the University of Heidelberg, and colleagues. As noted by the authors, the genetic map presented in the study offers a starting point for the design of antiviral agents targeting NS1, with the goal of suppressing viral replication as well as severe disease manifestations.

Dengue virus is one of the most prevalent mosquito-transmitted human pathogens. Despite the serious socio-[economic impact](#) of dengue-associated diseases, the only licensed vaccine has limited efficacy and an antiviral therapy is not available. NS1 is secreted from infected cells, counteracts antiviral immune responses, and contributes to the severe clinical manifestations of dengue. In addition, NS1 is essential for the [viral replication](#) cycle, but the underlying molecular mechanism is unknown. To address this gap in knowledge, Bartenschlager and colleagues used a combination of genetic, biochemical and imaging approaches to determine the role of NS1 in the [viral replication cycle](#).

The researchers identified a cluster of amino acid residues in NS1 that is important for efficient secretion of this protein. Moreover, they identified a novel interaction between NS1 and the precursor of a [viral protein](#) called NS4A-2K-4B; this interaction is required for viral RNA replication. The researchers also demonstrated that NS1 is required for

the generation of the membranous [dengue virus](#) replication organelle in infected cells. This function does not require RNA replication and is independent from NS1's interaction with NS4A-2K-4B. Taken together, the results provide new insights into the role of NS1 in viral RNA replication and establish a [genetic map](#) of residues in NS1 required for the diverse functions of this protein.

The authors add, "A detailed study of nonstructural protein 1 of Dengue virus identified determinants required for secretion of this protein, viral RNA replication and formation of the membranous replication organelle of this virus. A novel interaction of NS1 with a previously unrecognized NS4A-2K-NS4B precursor of the virus was found to be required for Dengue virus replication."

More information: A novel interaction between dengue virus nonstructural protein 1 and the NS4A-2K-4B precursor is required for viral RNA replication but not for formation of the membranous replication organelle. *PLoS Pathog* 15(5): e1007736.
doi.org/10.1371/journal.ppat.1007736

Provided by Public Library of Science

Citation: How the dengue virus replicates in infected cells (2019, May 9) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-05-dengue-virus-replicates-infected-cells.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
