

Diabetic amputations a 'shameful metric' of inadequate care

May 3 2019, by Anna Gorman, Kaiser Health News



Credit: CC0 Public Domain

On his regular rounds at the University of Southern California's Keck Hospital, Dr. David Armstrong lives a brutal injustice of American health care.

Each week, dozens of patients with diabetes come to him with deep wounds, severe infections and poor circulation—debilitating complications of a disease that has spiraled out of control. He works to save their limbs, but sometimes Armstrong and his team must resort to amputation to save the patient, a painful and life-altering measure he knows is nearly always preventable.

For decades now, the American medical establishment has known how to manage diabetes. Even as the number of people living with the illness continues to climb—today, estimated at more than 30 million nationwide—the prognosis for those with access to good health care has become far less dire. With the right medication, diet and lifestyle changes, patients can learn to manage their diabetes and lead robust lives.

Yet across the country, surgeons still perform tens of thousands of diabetic amputations each year. It's a drastic procedure that stands as a powerful example of the consequences of being poor, uninsured and cut off from a routine system of quality health care.

"Amputations are an unnecessary consequence of this devastating disease," said Armstrong, professor of surgery at Keck School of Medicine of USC. "It's an epidemic within an epidemic. And it's a problem that's totally ignored."

In California, where doctors performed more than 82,000 diabetic amputations from 2011 to 2017, people who were black or Latino were more than twice as likely as non-Hispanic whites to undergo amputations related to diabetes, a Kaiser Health News analysis found.

The pattern is not unique to California. Across the country, studies have shown that diabetic amputations vary significantly not just by race and ethnicity but also by income and geography. Diabetic patients living in

communities that rank in the nation's bottom quartile by income were nearly 39% more likely to undergo major amputations compared with people living in the highest-income communities, according to one 2015 study.

A 2014 study by UCLA researchers found that people with diabetes in poorer neighborhoods in Los Angeles County were twice as likely to have a foot or leg amputated than those in wealthier areas. The difference was more than tenfold in some parts of the county.

Amputations are considered a "mega-disparity" and dwarf nearly every other health disparity by race and ethnicity, said Dr. Dean Schillinger, a medical professor at the University of California-San Francisco. To begin with, people who are black or Latino are more at risk of diabetes than other groups—a disparity often attributed to socioeconomic factors such as higher rates of poverty and lower levels of education. They also may live in environments with less access to healthy food or places to exercise.

Then, among those with the disease, blacks and Latinos often get diagnosed after the disease has taken hold and have more complications, such as amputations. "If you go into low-income African American neighborhoods, it is a war zone," said Schillinger, former chief of the Diabetes Prevention and Control Program at the California Department of Public Health. "You see people wheeling themselves around in wheelchairs."

Part of the outrage for researchers is that medical science has made so much headway in diabetes treatment. Nationwide, fewer than 5 adults out of every 1,000 with diabetes get amputations.

But for those who do, the consequences are profound. More than half of amputations in California from 2011 to 2017 occurred among people

ages 45 to 64, according to the KHN analysis, meaning many people are left disabled and dependent on others for care during their prime working years.

Jackson Moss leaned back on his couch and raised his right leg. His wife, Bernadette, sprayed antiseptic on a gaping wound on the sole of his foot before dabbing it with Vaseline and rewrapping it with gauze.

A stocky man who used to deliver poultry, Moss, 47, said he had to stop working after his left leg was amputated below the knee about 10 years ago. Later, he lost part of his right foot. With Bernadette's help, he is trying to save the rest of it.

"If I didn't have my wife, I don't know where I'd be," said Moss, who wears a prosthesis on his left leg and uses a wheelchair. "I can't get around good like I used to."

Moss, who lives in Compton, embodies many of the characteristics of people most likely to get diabetic amputations. He is African American with a relatively low family income: about \$30,000 a year from his Social Security disability check and his wife's job with the county mental health department.

Moss has not always received regular medical care. His mother, who also had a leg amputated from diabetes, would take him to the doctor when he was a boy. But he stopped going as an adult. He didn't have insurance during much of his 20s and 30s. Medical care just wasn't a priority, he said, until about 25 years ago when his blood sugar shot up so high he passed out at home.

After he was diagnosed with Type 2 diabetes, he started seeing a physician more often. He tried to avoid sugar, as his doctor recommended, but bad habits die hard. "It takes a lot to eat right," he

said, "and it costs more."

One day, about 10 years ago, he bumped his toe on the bed. He thought little of it until he developed an infected wound. A fever sent him to the hospital, where his lower leg was removed. A few years later, with his diabetes still poorly controlled, he lost the toes on his other foot.

In recent years, Moss and his wife said, health providers have sometimes ignored their concerns. They recalled trips to the emergency room when they had to convince doctors his fever came from a diabetes-related infection. "They wouldn't take my word," he said. The couple did not see it as discrimination, more like dismissiveness.

Now, Moss goes to a clinic run by Martin Luther King, Jr. Community Hospital, which serves a large Latino and black population in South Los Angeles. On a recent visit, his doctor asked if he was staying off the foot with the wound. "I just get up when I have to go to the restroom and to get in and out of the bed," Moss responded.

Moss hopes someday he will be able to do more—get back to taking his grandsons to Chuck E. Cheese or playing dominoes with friends.

"I just sit here all day long," he said.

Amputations typically start with poorly controlled diabetes, a disease characterized by excess sugar in the blood. Untreated, it can lead to serious complications such as kidney failure and blindness.

People with diabetes often have reduced sensation in their feet, as well as poor circulation. As many as one-third of people with the most common form—Type 2—develop foot ulcers or a break in the skin that can become infected.

Amputations occur after those infections rage out of control and enter the bloodstream or seep deeper into the tissue. People with diabetes often have a condition that makes it harder for blood to circulate and wounds to heal.

The circumstances that give rise to amputations are complex and often intertwined: Patients may avoid doctors because their family and friends do, or clinics are too far away. Some may delay medical visits because they don't trust doctors or have limited insurance. Even when they seek treatment, some find it difficult to take medication as directed, adhere to dietary restrictions or stay off an infected foot.

Californians with diabetes who have a regular place to go for health care other than the emergency room are less likely to get amputations, according to an analysis conducted for Kaiser Health News by the UCLA Center for Health Policy Research. If they have a plan to control their diabetes, they also have less chance of amputation.

The analysis shows that many amputations could be avoided with better access to care and better disease management, said Ninez Ponce, director of the center.

"It's the most shameful metric we have on quality of care," Ponce said. "It is a health equity issue. We are a very rich state. We shouldn't be seeing these diabetic amputations."

An amputation often leads to a cascade of setbacks: more infections, more amputations, decreased mobility, social isolation. Research shows as many as three-quarters of people with diabetes who have had lower-limb amputations die within five years.

The health system bears surprisingly large costs for what remains a relatively uncommon problem. A single lower-limb [amputation](#) can cost

more than \$100,000. By far, government programs—Medicaid and Medicare—pay for the most amputations.

Experts say the best bet is to intervene well before they become necessary. People with diabetes are "very much in need of the simplest, basic, cost-effective, easy-to-implement treatments," said Dr. Philip Goodney, director of the Center for the Evaluation of Surgical Care at Dartmouth.

Along with basic measures to control diabetes, regular foot exams are key. The Centers for Disease Control and Prevention estimates somewhere between 11% and 28% of people with diabetes get the recommended podiatric care, a yearly foot exam to check for loss of sensation and blood flow. Under federal rules governing Medicaid, the government program for low-income Americans, such care is optional and not covered by every state.

California includes it as an optional benefit, limiting access to such care. An analysis by UCLA researchers estimated that the use of preventive podiatric services saved the Medi-Cal system—California's version of Medicaid—up to \$97 million in 2014, based on avoided hospital admissions and amputations, and that savings could be much greater if more patients had access.

Jesse Guerrero is 12, but already knows what diabetes—and amputations—can do to a family. He has seen how life changed since his mom, Patricia Zamora, had her first surgery. She had to stop working as a group home supervisor. They were evicted and eventually moved into his grandmother's house in Pomona.

Now, they stay home a lot more than they used to. "I want her to get better so we can finally go places," Jesse said.

First diagnosed with gestational diabetes, Zamora, 49, eventually was diagnosed with conventional Type 2. Though her mother has diabetes, she said, she didn't understand the risks.

Her serious troubles started in 2014, when she stubbed her big toe and it turned black and purple. When she finally went to an ER, doctors said it had to be amputated. The next year, after another stumble and another infection, doctors removed the remaining toes on her right foot.

Now, she is fighting a third wound and risks losing the limb below her knee. She uses a scooter and wears a boot to keep the pressure off.

Many days, she wants to give up.

"But I can't," she said. "I have Jesse."

His health is also a concern. Though only in middle school, Jesse is overweight, putting him at greater risk for Type 2 diabetes. She recently took away his PlayStation and signed him up for flag football so he would be more active.

Jesse, too, is scared.

"I don't want to get my foot cut off," he said. "I'd rather have a full life than a short one."

As hospitals have seen the impacts—and cost—of amputations, some have made efforts to reduce them. Some, like Keck Hospital, have started limb preservation centers, which use cross-disciplinary teams and technology to treat wounds and help patients improve disease management.

Even with a team of specialists, however, saving a limb often depends on

patients coming in early rather than waiting until their foot has become dangerously infected. But because their sensation is dulled, they often don't appreciate the danger.

"How do you get someone to come in if they don't have pain?" Armstrong said. "They need the gift of pain."

One of Armstrong's patients, Cirilo Delgado, has a wound on his heel that could cost him his lower leg. He already lost a toe.

Delgado, 41, knew diabetes ran in his family. His father, 68, has diabetes. His mother, who had diabetes and kidney failure, died at 67. His diabetic sister died at 35 of a heart attack, a possible complication of diabetes.

"I saw them die young," he said. "I don't want to be the next one."

Like Moss, Delgado didn't always have insurance. And he didn't seek care for his [diabetes](#) until the symptoms got dire.

Delgado used to work at a dry cleaning shop but had to stop because he doesn't have the balance he once did. His blood pressure fluctuates dangerously, and he needs dialysis three times a week for kidney failure. He has moved in with his father, a truck driver who stopped working to help care for him.

In November, doctors used a skin flap from his leg to try to heal his latest wound. He's praying he doesn't get another.

"I know there's a prosthesis," he said, "but it's not the same as a limb."

©2019 Kaiser Health News

Distributed by Tribune Content Agency, LLC.

Citation: Diabetic amputations a 'shameful metric' of inadequate care (2019, May 3) retrieved 22 March 2023 from

<https://medicalxpress.com/news/2019-05-diabetic-amputations-shameful-metric-inadequate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.