

New evidence links ultra-processed foods with a range of health risks

May 29 2019



Credit: CC0 Public Domain

Two large European studies published by *The BMJ* today find positive associations between consumption of highly processed ("ultra-processed") foods and risk of cardiovascular disease and death.

The researchers say further work is needed to better understand these effects, and a direct (causal) link remains to be established, but they call for policies that promote consumption of fresh or minimally processed foods over highly processed foods.

Ultra-processed foods include packaged baked goods and snacks, fizzy drinks, sugary cereals, ready meals containing food additives, dehydrated vegetable soups, and reconstituted meat and fish products—often containing high levels of added sugar, fat, and/or salt, but lacking in vitamins and fibre. They are thought to account for around 25-60% of daily energy intake in many countries.

Previous studies have linked ultra-processed foods to higher risks of obesity, [high blood pressure](#), high cholesterol, and some cancers, but firm evidence is still scarce.

In the first study, researchers based in France and Brazil assessed potential associations between ultra-processed foods and risk of cardiovascular and cerebrovascular disease (conditions affecting [blood supply](#) to the heart and brain).

Their findings are based on 105,159 French adults (21% men; 79% women) with an average age of 43 years who completed an average of six 24-hour dietary questionnaires to measure usual intake of 3,300 different food items, as part of the NutriNet-Santé study.

Foods were grouped according to degree of processing and rates of disease were measured over a maximum follow-up of 10 years (2009-2018).

Results showed that an absolute 10% increase in the proportion of ultra-processed food in the diet was associated with significantly higher rates of overall cardiovascular disease, coronary heart disease, and

[cerebrovascular disease](#) (increase of 12%, 13%, and 11% respectively).

In contrast, the researchers found a significant association between unprocessed or minimally processed foods and lower risks of all reported diseases.

In the second study, researchers based in Spain evaluated possible associations between ultra-processed food intake and risk of death from any cause ("all cause mortality").

Their findings are based on 19,899 Spanish university graduates (7,786 men; 12,113 women) with an average age of 38 years who completed a 136-item dietary questionnaire as part of the Seguimiento Universidad de Navarra (SUN) study.

Again, foods were grouped according to degree of processing and deaths were measured over an average of 10 years.

Results showed that higher consumption of ultra-processed foods (more than 4 servings per day) was associated with a 62% increased risk of all cause mortality compared with lower consumption (less than 2 servings per day). For each additional daily serving of ultra-processed food, mortality risk relatively increased by 18% (a dose-response effect).

Both studies are observational so can't establish causality, and there's a possibility that some of the observed risks may be due to unmeasured confounding factors.

Nevertheless, both studies took account of well known lifestyle risk factors and markers of dietary quality, and the findings back up other research linking highly processed food with [poor health](#).

As such, both research teams say policies that limit the proportion of

ultra-processed foods in the diet and promote consumption of unprocessed or minimally processed foods are needed to improve global public health.

This view is supported by Australian researchers in a linked editorial, who say the dietary advice is relatively straightforward: eat less ultra-processed food and more unprocessed or minimally processed food.

They say future research should explore associations between ultra-processed food and health harms in different populations around the world, and examine how harm occurs (for example by changing the gut microbiome in ways that could disturb energy balance).

In the meantime, policy makers "should shift their priorities away from food reformulation—which risks positioning ultra-processed [food](#) as a solution to dietary problems—towards a greater emphasis on promoting the availability, affordability, and accessibility of unprocessed or minimally processed foods," they conclude.

More information: Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Santé) *BMJ* (2019). www.bmj.com/content/365/bmj.11451

Association between consumption of ultra-processed foods and all cause mortality: SUN prospective cohort study, *BMJ* (2019). www.bmj.com/content/365/bmj.1949

Ultra-processed food and adverse health outcomes, *BMJ* (2019). www.bmj.com/content/365/bmj.12289

Provided by British Medical Journal

Citation: New evidence links ultra-processed foods with a range of health risks (2019, May 29) retrieved 13 May 2024 from <https://medicalxpress.com/news/2019-05-evidence-links-ultra-processed-foods-range.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.