

Fibro-adipose vascular anomaly: Old wine or new cocktail?

May 8 2019

Unique clinico-radiological features of a provisionally unclassified vascular anomaly can assist radiologists in identifying this uncommon distinct entity, according to a study to be presented at the ARRS 2019 Annual Meeting, set for May 5-10 in Honolulu, HI.

The study was conducted to evaluate the clinical and radiological features of fibro-adipose vascular anomaly (FAVA) including the use of advanced imaging techniques.

FAVA is a recently described soft-tissue vascular lesion, frequently misdiagnosed as venous malformation, mesenchymal tumor, or intramuscular hemangioma and refractory to conventional modalities of treatment. In a retrospective review of hospital records, the clinical and imaging findings of 20 patients were reviewed to construct a profile of the unique clinico-radiological features of FAVA.

The typical clinical profile is a young patient with painful intramuscular lesion in the extremities, causing contractures. The lesion is solid-appearing with heterogeneous echotexture on ultrasound, soft on elastography, and shows progressive heterogeneous enhancement on contrast ultrasound and dynamic magnetic resonance imaging (MRI) with associated phlebectasia.

"FAVA is a new and novel entity that the radiologists should be aware of," said Dr. Ankur Goyal, author of the study "as they may be first ones to suggest this possibility."

More information: www.arrs.org/am19

Provided by American Roentgen Ray Society

Citation: Fibro-adipose vascular anomaly: Old wine or new cocktail? (2019, May 8) retrieved 23 May 2024 from <https://medicalxpress.com/news/2019-05-fibro-adipose-vascular-anomaly-wine-cocktail.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.