

Glucosamine supplements may be linked to lower risk of cardiovascular disease

May 14 2019



Human heart. Credit: copyright American Heart Association

Regular use of glucosamine supplements may be related to a lower risk of cardiovascular disease (CVD) events, suggests an analysis of data from the UK Biobank study, published in *The BMJ* today.

The findings suggest that glucosamine may have benefits in preventing CVD events, such as coronary heart disease and stroke, and further clinical trials are needed to test this theory, say the researchers.

Glucosamine is a popular dietary supplement used to relieve osteoarthritis and joint pain. While its effectiveness on [joint pain](#) continues to be debated, emerging evidence suggests that glucosamine may have a role in preventing cardiovascular disease and reducing mortality. However, conclusive evidence is still lacking

So to explore these potential associations further, researchers led by Professor Lu Qi at Tulane University in New Orleans drew on data from the UK Biobank—a large population based study of more than half a million British men and women.

Their analysis included 466,039 participants without CVD, who completed a questionnaire on supplement use, including glucosamine.

Death certificates and hospital records were then used to monitor CVD events, including CVD death, [coronary heart disease](#) (CHD), and stroke, over an average seven-year follow up period.

Overall, almost one in five (19.3%) participants reported glucosamine use at the start of the study.

The researchers found that glucosamine use was associated with a 15% lower risk of total CVD events, and a 9% to 22% lower risk of CHD, stroke, and CVD death compared with no use.

These favourable associations remained after taking account of traditional risk factors, including age, sex, weight (BMI), ethnicity, lifestyle, diet, medication and other supplement use.

The association between glucosamine use and CHD was also stronger in current smokers (37% lower risk) compared with never (12%) and former smokers (18%).

Several mechanisms may explain these results, say the authors. For example, regular use of glucosamine has been linked to a reduction in levels of C-reactive protein (CRP), a chemical associated with inflammation. This may also help to explain the stronger association among smokers, who have higher levels of inflammation and higher risk of CVD than non-smokers.

In addition, previous data suggest that glucosamine may mimic a low carbohydrate diet, which has been inversely associated with the development of CVD.

Despite the large sample size, this is an observational study, and as such, can't establish cause, and the researchers point to some limitations, such as lack of information on dose, duration, and side effects of glucosamine use.

Regular [glucosamine](#) use may also be a marker for a [healthy lifestyle](#), they add, but this is unlikely to have affected the results.

As such, they conclude that "habitual use of [glucosamine](#) supplements to relieve osteoarthritis pain might also be related to lower risks of CVD events. Further [clinical trials](#) are warranted to test this hypothesis."

More information: Association of habitual glucosamine use with risk of cardiovascular disease: prospective study in UK Biobank, *BMJ* (2019). [DOI: 10.1136/bmj.11628](https://doi.org/10.1136/bmj.11628) , www.bmj.com/content/365/bmj.11628

Provided by British Medical Journal

Citation: Glucosamine supplements may be linked to lower risk of cardiovascular disease (2019, May 14) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2019-05-glucosamine-supplements-linked-cardiovascular-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.