

# GPs need training to tackle chronic opioid use

May 13 2019

---



Credit: CC0 Public Domain

GPs must be better-equipped to support patients to manage the psychological challenge of reducing their opioid use—according to new research from the University of East Anglia.

The recommendation is part of a toolkit being launched today to help GPs reduce the amount of opioids they prescribe.

The toolkit outlines seven areas of best practice to tackle chronic [opioid use](#)—based on international research evidence, the experiences of health organisations and individual practitioners.

It comes after figures for England and Wales revealed an increase in [opioid prescriptions](#) of more than 60 per cent over the last decade—from 14 million in 2008 to 23 million last year.

Lead researcher Dr. Debi Bhattacharya, from UEA's School of Pharmacy, said: "Opioids, like morphine, tramadol and fentanyl, can be effective for the short-term management of severe pain. However, they are highly addictive which makes stopping difficult yet long-term use can impair quality of life and overuse can be deadly."

"GPs and other [health professionals](#) need to urgently, proactively work with patients prescribed long-term opioids for non-cancer pain to gradually reduce or 'taper' their doses.

"But if GPs are expected to initiate discussions about tapering or stopping opioids, they must be equipped with training to manage the psychological challenges experienced by patients when trying to reduce their opioid use."

"Without this training, prescribers are reticent to open 'a can of worms' that they know they don't have the skills to manage."

"For opioid tapering interventions to be effective, GPs need training in giving their patients the skills to manage any withdrawal effects."

Dr. Bhattacharya said: "There needs to be a clear expectation that opioid

de-prescribing is the responsibility of the prescriber.

"Incentives may help GPs and other health professionals to prioritise reducing the amount of opioids being prescribed to [patients](#), particularly among those who have been taking them long-term.

"Prescribers need to better understand the consequences of excess [opioid](#) use, and they need better guidelines about how to gradually reduce or 'taper' doses."

**More information:** The 'Toolkit for Tackling Chronic Opioid Use in Non-Cancer Pain' launches on Monday May 13, and is available at [www.uea.ac.uk/pharmacy/research...-cancer-pain/toolkit](http://www.uea.ac.uk/pharmacy/research/2019-05-gps-tackle-chronic-opioid.html)

Provided by University of East Anglia

Citation: GPs need training to tackle chronic opioid use (2019, May 13) retrieved 2 May 2024 from <https://medicalxpress.com/news/2019-05-gps-tackle-chronic-opioid.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.