

Novel index accurately predicts PCI success post procedure compared to standard measurement metrics

May 21 2019

Results from a comprehensive analysis demonstrate the effectiveness of measuring a non-hyperemic pressure ratio (NHPR), pressure distal/pressure aortic (Pd/Pa) alongside fractional flow reserve (FFR) post percutaneous coronary intervention (PCI). This prospective study validates the diagnostic accuracy of Pd/Pa in identifying residual ischemia post intervention against the reference standard, FFR. The study is scheduled to be presented on Wednesday, May 22 at the Society for Cardiovascular Angiography and Interventions (SCAI) 2019 Scientific Sessions.

PCI is a [nonsurgical procedure](#) that improves [blood flow](#) to the heart by using a stent to open up [blood vessels](#). Since 1998, more than 17.5 million [coronary angiography](#) and PCI procedures have been performed in the United States (*Journal of the American College of Cardiology*).

While the value of FFR (a way to measure coronary pressure drop across a stenosis; synonymous to flow under maximal hyperemia) to determine appropriateness of revascularization is well-established and incorporated into current guidelines, there is a growing body of evidence supporting the role of remeasuring FFR after [intervention](#). It has been shown that almost 20-30 percent of [patients](#) continue to have residual ischemia (suboptimal flow) after successful stenting as determined by angiography- the current standard for determining procedural success. Furthermore, post-PCI FFR has been associated with long term cardiac

outcomes in multiple studies including post hoc analyses of the pivotal FAME trials.

Despite the robust evidence base, use of FFR in general is very low, especially post-PCI. Multiple factors including additional procedure time, cost and patient side effects have been cited as potential reasons. Study authors sought to build on this body by evaluating and validating the diagnostic performance of a non-hyperemic index (Pd/Pa), universally available on all [pressure](#) wire platforms, against the current "gold standard" of FFR post-PCI.

The study evaluated more than 1,200 patients and 1,600 vessels comparing accuracy of Pd/Pa pre-and-post PCI against FFR. The diagnostic accuracy and comparative performance of Pd/Pa was then prospectively validated in 230 patients and 264 vessels against the reference standard FFR in identifying ischemia (

Citation: Novel index accurately predicts PCI success post procedure compared to standard measurement metrics (2019, May 21) retrieved 29 June 2024 from <https://medicalxpress.com/news/2019-05-index-accurately-pci-success-procedure.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
