

Prehospital Tx guidelines may improve outcomes in severe TBI

May 10 2019



(HealthDay)—Implementation of prehospital traumatic brain injury

(TBI) guidelines is associated with improvement in survival to hospital discharge for patients with severe TBI, according to a study published online May 8 in *JAMA Surgery*.

Daniel W. Spaite, M.D., from the University of Arizona in Phoenix, and colleagues conducted an intention-to-treat study using a before/after controlled design with patients with moderate to critically severe TBI. The correlation between implementing the nationally vetted, evidence-based prehospital treatment [guidelines](#) and outcomes was examined. A total of 21,852 patients met inclusion criteria for analysis.

The researchers found that the adjusted odds ratio (aOR) was 1.06 (95 percent confidence interval [CI], 0.93 to 1.21; $P = 0.40$) for survival to [hospital discharge](#) in the primary analysis. For survival to hospital admission, the aOR was 1.70 (95 percent CI, 1.38 to 2.09; P survival to discharge).

"We found a therapeutic sweet spot and showed that the guidelines had an enormous impact on people with severe TBI," Spaite said in a statement. "The guidelines did not make a difference in the moderate TBI group because those individuals would most likely have survived anyway and, unfortunately, the extent of injuries sustained in many critical [patients](#) was too extreme to overcome."

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Prehospital Tx guidelines may improve outcomes in severe TBI (2019, May 10) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-05-prehospital-tx-guidelines-outcomes-severe.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.