


Research reveals possible strategy for blocking the spread of pneumococcal disease

May 23 2019



Rosch and his colleagues used high-throughput genetic screening of more than 65,000 *S. pneumoniae* mutants to search for factors required for transmission. Credit: St. Jude Children's Research Hospital

St. Jude Children's Research Hospital scientists have reduced transmission of pneumococci in mice using an experimental vaccine that

targets proteins on the surface of the bacteria. The research appears online today in the journal *Cell Host & Microbe*.

The results suggest a possible strategy for eliminating *Streptococcus pneumoniae* (pneumococcus) with combination vaccines that block bacterial transmission and prevent invasive disease. Current vaccines have helped reduce pneumococcal disease, but have done little to stop bacterial transmission.

S. pneumoniae is the most common cause of sepsis, pneumonia, meningitis and middle ear infections in [young children](#). Cancer patients, people 65 and older and others are also at an increased risk of invasive disease.

"A key aspect of pneumococci's success is the ability to transmit from one host to another, yet that process and the factors involved are not well understood," said corresponding author Jason Rosch, Ph.D., an associate member of the St. Jude Department of Infectious Diseases. "These data suggest that rationally designed [combination vaccines](#) may offer a way to disrupt the spread and eliminate invasive disease that threatens young and old alike."

S. pneumoniae lives in the nasal passages of most people at some point in their life, often beginning at an early age. Once pneumococcus has colonized a host, the germ can infect others or move deeper into the body and cause invasive disease. Vaccines are available to prevent invasive disease, but they are not as effective at blocking transmission.

Transmission essentials

Rosch and his colleagues used high-throughput genetic screening of more than 65,000 *S. pneumoniae* mutants to search for factors required for transmission.

Using a model developed by St. Jude biostatisticians, the researchers identified about 70 bacterial proteins that played an important role in transmission from infected to uninfected ferrets. Investigators confirmed the findings in mice.

Many of the proteins were involved in metabolism, [transcriptional regulation](#) and competency (bacterial exchange of genetic material).

The transmission factors also included surface proteins—PiaA and CppA. Vaccinating female mice with either or both proteins blocked transmission of pneumococcus to their offspring compared to those born to mice who received a current vaccine.

More information: Hannah M. Rowe et al. Bacterial Factors Required for Transmission of *Streptococcus pneumoniae* in Mammalian Hosts, *Cell Host & Microbe* (2019). [DOI: 10.1016/j.chom.2019.04.012](https://doi.org/10.1016/j.chom.2019.04.012)

Provided by St. Jude Children's Research Hospital

Citation: Research reveals possible strategy for blocking the spread of pneumococcal disease (2019, May 23) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-05-reveals-strategy-blocking-pneumococcal-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--