


Walking and strength training may decrease the risk of dying from liver disease

May 20 2019



Credit: CC0 Public Domain

Physical activity, including walking and muscle-strengthening activities, were associated with significantly reduced risk of cirrhosis-related death, according to research presented at Digestive Disease Week (DDW) 2019. Chronic liver disease is increasing, partly due to the obesity epidemic, and currently there are no guidelines for the optimal type of

exercise for the prevention of cirrhosis-related mortality. Researchers hope these findings will help provide specific exercise recommendations for patients at risk for cirrhosis and its complications.

"The benefit of exercise is not a new concept, but the impact of exercise on mortality from cirrhosis and from [liver cancer](#) has not yet been explored on this scale," said Tracey Simon, MD, lead researcher on the study and instructor of medicine at Harvard Medical School and Massachusetts General Hospital, Boston. "Our findings show that both walking and [strength training](#) contribute to substantial reductions in risk of cirrhosis-related death, which is significant because we know very little about modifiable risk factors."

Dr. Simon and her team prospectively followed 68,449 women from the Nurses' Health Study and 48,748 men from the Health Professionals Follow-up Study, without known [liver disease](#) at baseline. Participants provided highly [accurate data](#) on physical activity, including type and intensity, every two years from 1986 through 2012, which allowed researchers to prospectively examine the association between physical activity and cirrhosis-related death.

Researchers observed that adults in the highest quintile of weekly walking activity had 73 percent lower risk for cirrhosis-related death than those in the lowest quintile. Further risk reduction was observed with combined walking and muscle-strengthening exercises.

Previous research has been limited to studies that assessed physical activity at just one point in time, or studies with very short-term follow-up. This was the first prospective study in a large U.S. population to include detailed and updated measurements of physical activity over such a prolonged period, which allowed researchers to more precisely estimate the relationship between physical activity and liver-related outcomes.

"In the U.S., mortality due to cirrhosis is increasing dramatically, with rates expected to triple by the year 2030. In the face of this alarming trend, information on modifiable risk factors that might prevent liver disease is needed," said Dr. Simon. "Our findings support further research to define the optimal type and intensity of [physical activity](#) to prevent adverse outcomes in patients at risk for cirrhosis."

Provided by Digestive Disease Week

Citation: Walking and strength training may decrease the risk of dying from liver disease (2019, May 20) retrieved 9 April 2024 from <https://medicalxpress.com/news/2019-05-strength-decrease-dying-liver-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--