

Risk of suicide attempt by children doubles if parent uses opioids

May 22 2019



Credit: CC0 Public Domain

The suicide rate among young people in the U.S. has risen dramatically in the past 15 years. Over the same time period, opioid use and abuse in adults also has increased considerably.

Researchers from the University of Chicago and the University of Pittsburgh reported on a connection between these two epidemics in a



study published this week in JAMA Psychiatry.

"Until now, there has been little focus on the association between the increase in <u>opioid use</u> among adults and the risk of <u>suicidal behavior</u> by their children," said Robert D. Gibbons, Ph.D., Blum-Riese Professor of Biostatistics and director of the Center for Health Statistics at UChicago and senior author of the paper. "We theorized such a link was plausible because parental substance abuse is a known risk factor for suicide attempts by their children. In addition, depression and suicide attempts by <u>parents</u>—which are known to be related to suicidal behavior in their offspring—are more common among adults who abuse opioids."

The researchers analyzed data from more than 240,000 parents, ages 30 to 50, between 2010 and 2016. Half of the group had filled <u>opioid</u> prescriptions for at least 365 days. The other half had no history of using the drug during that time. The two groups were matched on a number of factors related to suicide attempts and opioid use. Rates of suicide attempts were studied in over 330,000 children, ages 10 to 19, from these two groups of parents over the same six-year period.

Of the children whose parents used opioids, 678 (0.37%) attempted suicide. Of the sons and daughters of parents who did not use opioids, 212 (0.14%) made a suicide attempt. The researchers found that opioid use by a parent is associated with a doubling of the risk of <u>suicide</u> attempts by their children. The results were statistically significant even when adjusted for child age and sex, depression or <u>substance use</u> <u>disorder</u> in child or parent, and history of a suicide attempt in a parent.

"These findings demonstrate that opioid use by a parent or parents doubles the risk for suicidal behavior by their children," said David A. Brent, MD, psychiatrist and chair of suicide studies at the University of Pittsburgh, also an author on the paper. "The epidemics of adult opiate abuse and child suicidal behavior appear to be linked, and the disturbing



upward trends in mortality due to opiates and due to child suicide may have common roots."

Gibbons and Brent call for improved diagnosis and treatment of parents who use opioids as well as mental health screening and referral to care for their <u>children</u>. "These actions could help reverse the upward trend in deaths due to the twin epidemics of <u>suicide</u> and opioid overdose," Gibbons said.

More information: "Association Between Parental Medical Claims for Opioid Prescriptions and Risk of Suicide Attempt by Their Children" *JAMA Psychiatry* (2019). <u>DOI:</u> <u>10.1001/jamapsychiatry.2019.0940</u>

Provided by University of Chicago Medical Center

Citation: Risk of suicide attempt by children doubles if parent uses opioids (2019, May 22) retrieved 20 April 2024 from <u>https://medicalxpress.com/news/2019-05-suicide-children-parent-opioids.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.