

Weight-loss surgery boosts success of procedure to fix A-fib

May 8 2019, by Steven Reinberg, Healthday Reporter



(HealthDay)—Profoundly obese people are prone to an irregular

heartbeat known as atrial fibrillation, but new research suggests that weight-loss surgery can improve the odds that a procedure to restore a normal heart rhythm will work.

When [atrial fibrillation](#), or a-fib, cannot be controlled by medications, a procedure called ablation can help. It targets the nerves in the heart where atrial fibrillation occurs. By electrically burning these areas, the heartbeat may return to normal.

"Bariatric [weight-loss] [surgery](#) before ablation was associated with a threefold reduction in atrial fibrillation recurrence rates," said researcher Dr. Eoin Donnellan, a fellow in cardiology at the Cleveland Clinic.

Both obesity and a-fib are at epidemic proportions in the United States, and both can increase the risk of cardiovascular disease, the researchers said.

As many as one in five cases of a-fib are related to obesity, and obesity has been identified as a risk factor for the heart condition, the researchers noted.

Some patients who have weight-loss surgery may see their a-fib stop on its own or can have it controlled with medication, Donnellan said. But for many, ablation may be the best way to cure it.

Although ablation has become a common procedure to treat a-fib, obese patients are more likely to have their abnormal heartbeat return after the procedure, he said.

"While previously, morbidly obese patients with atrial fibrillation that can't be controlled with drugs were considered ineligible for ablation, [bariatric surgery](#) offers them an option to get rid of the atrial fibrillation," Donnellan explained.

For this study, Donnellan and his colleagues collected data on nearly 240 obese patients undergoing ablation for atrial fibrillation.

Among these patients, 51 had weight-loss surgery before ablation. The researchers then looked for recurrence rates of a-fib in the three years following the procedure.

The investigators found that among patients who didn't have weight-loss surgery, almost 61% had a recurrence of their a-fib, compared to just under 20% of the patients who had weight-loss surgery before ablation.

In addition, only 12% of those who had weight-loss surgery needed another ablation, compared with 41% of those who hadn't had the surgery.

Losing weight has been associated with making atrial fibrillation easier to control, said Dr. Vivek Reddy. He is director of cardiac electrophysiology at the Icahn School of Medicine at Mount Sinai, in New York City.

Obesity directly affects heart tissue, making it more susceptible to a-fib, Reddy said.

But to prove whether [weight-loss surgery](#) really makes ablation more effective, a larger study that randomly assigns [obese patients](#) to have or not have surgery before [ablation](#) is needed, he explained.

This study, which was observational only, suggests that the surgery has a meaningful effect, he said. In any case, losing weight makes a-fib easier to treat.

"Lose weight however you can," Reddy said. "If you're morbidly obese and undergo bariatric surgery, there are benefits throughout the body,

including atrial fibrillation reduction."

The findings were scheduled for presentation Friday at the Heart Rhythm Society annual meeting, in San Francisco. Research presented at meetings should be considered preliminary until published in a peer-reviewed journal.

More information: For more about a-fib, visit the [American Heart Association](#).

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Weight-loss surgery boosts success of procedure to fix A-fib (2019, May 8) retrieved 6 May 2024 from <https://medicalxpress.com/news/2019-05-weight-loss-surgery-boosts-success-procedure.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--