

# Fewer than 60% of young women diagnosed with STIs in emergency departments fill scripts

May 28 2019

---



Monika K. Goyal, M.D., MSCE, assistant chief of Children's Division of Emergency Medicine and Trauma Services and the study's senior author. Credit: Children's National

Fewer than 60% of young women diagnosed with sexually transmitted infections (STIs) in the emergency department fill prescriptions for

antimicrobial therapy to treat these conditions, according to a research letter published online May 28, 2019, by *JAMA Pediatrics*.

Adolescents make up nearly half of the people diagnosed with sexually transmitted infections each year. According to the Centers for Disease Control and Prevention, untreated sexually transmitted diseases in women can cause [pelvic inflammatory disease](#) (PID), an infection of the reproductive organs that can complicate getting pregnant in the future.

"We were astonished to find that teenagers' rates of filling STI prescriptions were so low," says Monika K. Goyal, M.D., MSCE, assistant chief of Children's Division of Emergency Medicine and Trauma Services and the study's senior author. "Our findings demonstrate the imperative need to identify innovative methods to improve treatment adherence for this high-risk population."

The retrospective cohort study, conducted at two emergency departments affiliated with a large, urban, tertiary care children's hospital, enrolled adolescents aged 13 to 19 who were prescribed antimicrobial treatment from Jan. 1, 2016, to Dec. 31, 2017, after being diagnosed with PID or testing positive for chlamydia.

Of 696 emergency department visits for diagnosed STIs, 208 teenagers received outpatient prescriptions for antimicrobial treatments. Only 54.1% of those prescriptions were filled.

"Teenagers may face a number of hurdles when it comes to STI treatment, including out-of-pocket cost, access to transportation and confidentiality concerns," Dr. Goyal adds.

Future studies will attempt to identify barriers to filling prescriptions in order to inform development of targeted interventions based in the [emergency department](#) that promote adherence to STI treatment.

**More information:** Alexandra Lieberman et al, Frequency of Prescription Filling Among Adolescents Prescribed Treatment for Sexually Transmitted Infections in the Emergency Department, *JAMA Pediatrics* (2019). [DOI: 10.1001/jamapediatrics.2019.1263](https://doi.org/10.1001/jamapediatrics.2019.1263)

Provided by Children's National Medical Center

Citation: Fewer than 60% of young women diagnosed with STIs in emergency departments fill scripts (2019, May 28) retrieved 19 April 2024 from <https://medicalxpress.com/news/2019-05-young-women-stis-emergency-departments.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.