

# What influences critical care doctors in withdrawing life support for patients with brain injury?

June 17 2019

---



Credit: CC0 Public Domain

Decisions to withdraw life support treatments in critically ill patients with severe brain injury are complicated, are based on many factors, and

are usually made by critical care physicians and families in the intensive care unit. A study in *CMAJ* (*Canadian Medical Association Journal*) [provides](#) new understanding on the decision-making processes for this patient population, which accounts for most hospital deaths from trauma.

"Many clinicians struggle to make recommendations to withdraw life-sustaining treatments because decision-making is often complicated by uncertainty from trying to match family or caregiver opinions about what they think the patient would have wanted in terms of quality of life and how well physicians can predict a prognosis," says Dr. Alexis Turgeon, a critical care [physician](#) and researcher at CHU de Québec—Université Laval Research Centre and Canada Research Chair in Critical Care Neurology and Trauma at Université Laval, Québec City, Quebec.

Most deaths in critically ill patients with severe traumatic brain injury occur after a decision to withdraw life-sustaining treatments. This patient population differs from the general [intensive care unit](#) (ICU) population as most patients were healthy before admission to the ICU, as compared with [older patients](#) who may already have poorer quality of life due to pre-existing illness. Therefore, decisions to withdraw life-sustaining treatments are made differently—mainly based on long-term prognosis and quality of life.

Researchers performed a descriptive qualitative study of interviews with critical care physicians from across Canada to understand the factors that determine a critical care physician's decision to discuss with families the withdrawal of life-sustaining treatments in patients with severe traumatic brain injury. Results show that several factors are involved, including the patient's pre-expressed wishes and the family's wishes, severity and location of the injury, along with evidence. Past physician experience, legislation, opinions of colleagues and time are additional factors

influencing decisions. The incidence of withdrawal of life-sustaining treatments and of death in critically ill patients with traumatic brain injury varies between hospitals.

"A major factor for physicians in decision-making is the influence of the patient circumstances and the family," says Dr. Turgeon. "This is reassuring but challenging when the patient's prognosis is uncertain."

Better evidence, tools to help predict patient outcomes, standardization, better ways to integrate patient values and preferences into decision-making, improved training during critical care fellowships and more time to estimate prognosis are some things that could improve [decision-making](#) regarding whether to withdraw life-sustaining treatments.

"Our study has implications for the care of critically ill [patients](#) with traumatic brain injury, and we hope it will inform policy to improve how critical care physicians determine prognosis and level of care decisions with families," says Dr. Turgeon.

**More information:** *Canadian Medical Association Journal* (2019).  
[www.cmaj.ca/lookup/doi/10.1503/cmaj.190154](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.190154)

Provided by Canadian Medical Association Journal

Citation: What influences critical care doctors in withdrawing life support for patients with brain injury? (2019, June 17) retrieved 27 April 2024 from  
<https://medicalxpress.com/news/2019-06-critical-doctors-life-patients-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--