

Settling the debate on serotonin's role in sleep

June 24 2019



Two zebrafish brains, side by side, with serotonin molecules shown in red. The brain on the right comes from a fish that cannot synthesize serotonin in its raphe nuclei, leading to less of the neurochemical throughout the brain. This animal slept about half as much as a normal zebrafish (brain on the left). Credit: Caltech



Serotonin is a multipurpose molecule found throughout the brain, playing a role in memory, cognition, and feelings of happiness and other emotions. In particular, researchers have long debated serotonin's role in sleep: Does serotonin promote sleep, or its opposite, wakefulness?

Now, Caltech scientists have found that <u>serotonin</u> is necessary for sleep in <u>zebrafish</u> and mouse models.

A paper describing the research appears online on June 24 in the journal *Neuron*. The work is a collaboration between the Caltech laboratories of David Prober, professor of biology and affiliated faculty member of the Tianqiao and Chrissy Chen Institute for Neuroscience at Caltech; and Viviana Gradinaru (BS '05), professor of neuroscience and <u>biological engineering</u>, Heritage Medical Research Institute Investigator, and director of the Chen Institute's Center for Molecular and Cellular Neuroscience.

Previous studies on serotonin and sleep have yielded conflicting results. Some research showed that serotonin promotes sleep, but other work showed that serotonin-producing neurons were most active and releasing the chemical during wakefulness.

In order to settle this debate, the Caltech team focused on a region called the raphe nuclei, which has the brain's main population of serotoninproducing (or serotonergic) neurons. The raphe are evolutionarily ancient structures found in the brain stem of a wide range of organisms from fish to humans, and they are responsible for both manufacturing and sending out serotonin to other brain regions.

Led by senior postdoctoral scholar Grigorios Oikonomou of the Prober lab, the research began using zebrafish, tiny transparent fish that are widely used as a model to study sleep. Like humans, <u>zebrafish larvae</u> are diurnal—meaning that their sleep occurs mostly at night.



First, the researchers genetically mutated zebrafish so that their raphe did not produce serotonin. These mutant fish, the team found, slept about half as much as normal fish. In another experiment, the researchers removed the raphe altogether, and these fish also slept much less than usual.

"This suggests that serotonin produced by the raphe is required for the fish to get normal amounts of sleep," says Oikonomou.



Ball-and-stick model of the serotonin molecule. Credit: Public Domain



In a third set of experiments, zebrafish were genetically modified so that their raphe could be activated by light. Shining a light on these fish put them to sleep, implying that activation of the raphe induces sleep. This effect requires serotonin, because activating the raphe in <u>fish</u> that do not synthesize serotonin had no effect on sleep.

The team from the Prober laboratory then collaborated with scientists in the Gradinaru laboratory to continue the serotonin studies in mice. Led by graduate student Michael Altermatt, the team examined serotonergic neurons in the mouse raphe and confirmed that they are indeed mostly active while the animals are awake and less active during sleep, in agreement with previous studies.

As Prober's lab did in zebrafish, the Gradinaru lab's team genetically removed the serotonergic neurons in the mouse raphe and found that the mice slept less than usual. Stimulating these neurons with light also put the mice to sleep but only when the light was administered at frequencies that are consistent with the naturally occurring baseline activity pattern of these neurons during wakefulness.

"There's an obvious paradox here: stimulating the neurons causes the animals to sleep, and yet the neurons are normally active during the day," says Altermatt.

Oikonomou explains: "There are two main factors that control sleep. One is the <u>circadian clock</u>—when it is light during the day, the body is awake, and when it gets dark, the body knows to sleep. The other factor is called homeostatic sleep pressure. When you wake up in the morning, you have just gotten rest, and so you're energetic. As the day goes on, you get tired and sleepy, so there is a building of pressure to sleep. If you don't sleep that night, your sleep pressure is even higher, and you are even more tired the next day even though it's light outside, and your circadian clock dictates that you should be awake."



"The theory is that, in order to sleep, you need to have high sleep pressure and the circadian clock needs to be aligned with the time of day—nighttime for diurnal creatures like us and daytime for nocturnal animals."

The researchers theorize that the firing of <u>neurons</u> in the raphe and their release of serotonin is a way for the brain to build up sleep pressure. Indeed, they found that zebrafish lacking serotonin as well as mice with ablated raphe show reduced sleep pressure.

While the studies were in animal models, the raphe region and its production of serotonin are similar in human brains. The research can contribute to explanations of some <u>sleep</u>-related side effects of common antidepressant drugs that increase serotonin levels in the brain.

The paper is titled "The Serotonergic Raphe Promote Sleep in Zebrafish and Mice."

More information: "The Serotonergic Raphe Promote Sleep in Zebrafish and Mice" *Neuron* (2019). DOI: 10.1016/j.neuron.2019.05.038

Provided by California Institute of Technology

Citation: Settling the debate on serotonin's role in sleep (2019, June 24) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2019-06-debate-serotonin-role.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.