

## Excess weight and body fat cause cardiovascular disease

June 14 2019

**UK Biobank** N = 367 703

**Mendelian randomization** 

82 SNPs

96 SNPs



## RISK



Associations of BMI and fat mass index with cardiovascular conditions. Credit: European Heart Journal and Professor Susanna Larsson

Excess weight and body fat cause a range of heart and blood vessel



diseases, according to the first study to investigate this using a method called Mendelian randomisation.

In particular, the study published in the *European Heart Journal* today, shows that as body mass index (BMI) and fat mass increase, so does the risk of aortic <u>valve</u> stenosis—a condition in which the valve controlling the flow of blood from the <u>heart</u> to the body's largest blood vessel, the aorta, narrows and fails to open fully.

Mendelian randomisation is a way of showing whether or not individual risk factors actually cause disease, rather than just being associated with it. It uses genetic variants that are already known to be associated with potential risk factors, such as BMI and body fat, as indirect indicators or "proxies" for these risk factors. This enables researchers to discover whether the risk factor is the cause of the disease (rather than the other way around), and reduces bias in results because genetic variants are determined at conception and cannot be affected by subsequent external or environmental factors, or by the development of disease.

The researchers, led by Susanna Larsson, associate professor and senior researcher at the Karolinska Institute, Stockholm, Sweden, studied 96 genetic variants associated with BMI and body fat mass to estimate their effect on 14 cardiovascular diseases in 367,703 participants of white-British descent in UK Biobank—a UK-based national and international resource containing data on 500,000 people, aged 40-69 years.

She said: "The causal association between BMI and fat mass and several heart and blood vessel diseases, in particular aortic valve stenosis, was unknown. Using Mendelian randomisation we found that higher BMI and fat mass are associated with an increased risk of aortic valve stenosis and most other cardiovascular diseases, suggesting that excess <u>body fat</u> is a cause of cardiovascular disease."



People who had genetic variants that predict higher BMI were at increased risk of aortic valve stenosis, heart failure, <u>deep vein</u> thrombosis, high blood pressure, <u>peripheral artery disease</u>, <u>coronary</u> <u>artery disease</u>, atrial fibrillation and pulmonary embolism. For every genetically-predicted 1kg/m2 increase in BMI, the increased risk ranged from 6% for pulmonary embolism to 13% for aortic valve stenosis. (Above a BMI that is considered 'healthy' (20-25 kg/m2) every 1 kg/m2 increase in BMI for someone who is 1.7 metres tall (5'7") corresponds to a weight gain of nearly 3 kg.)

The researchers also found that risk of cardiovascular diseases increased with the genetic variants predicting increases in fat mass. The greatest increased risk was also for aortic valve stenosis (46% increased risk), followed by ischaemic stroke, transient ischaemic attack, atrial fibrillation, heart failure, peripheral artery disease, deep vein thrombosis, high blood pressure and coronary artery disease.

The researchers stress that although these genetic variants can predispose people to be more likely to gain excess weight, the most important factors implicated in the development of cardiovascular disease are diet and physical activity.

Professor Larsson said: "Our genes can make us somewhat more predisposed to gain body weight but lifestyle factors, such as overeating and lack of physical activity, are the major determinants of overweight. A <u>healthy diet</u> is the cornerstone of cardiovascular disease prevention, and how much we eat should be limited to the amount of energy required to maintain a healthy body weight, which is a BMI of between 20 to 25 kg/m2. People who are predisposed to a higher BMI may need to work a bit harder to maintain a healthy weight."

The strengths of the study include the large numbers of people involved and the fact that they were of European descent, which reduces the



potential for bias from different populations. Potential limitations are that some genetic variants may be associated with more than one characteristic, that the number of cases were few for some diseases, and that there was a lack of information on the severity of aortic valve stenosis.

The damaged valve in aortic valve stenosis means that less blood leaves the heart and it has to work harder to pump enough <u>blood</u> out to circulate round the body. Blood can back up in other parts of the heart and sometimes the lungs. This can lead to shortness of breath, tiredness, fainting, chest pain and an irregular heart beat.

**More information:** Susanna C Larsson et al, Body mass index and body composition in relation to 14 cardiovascular conditions in UK Biobank: a Mendelian randomization study, *European Heart Journal* (2019). DOI: 10.1093/eurheartj/ehz388

Provided by European Society of Cardiology

Citation: Excess weight and body fat cause cardiovascular disease (2019, June 14) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2019-06-excess-weight-body-fat-</u> <u>cardiovascular.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.