

# 'Goldilocks' neurons promote REM sleep

June 19 2019

---



Using a technique called optogenetics, a laser light was used to turn off or on MCH neurons time locked with the temperature warming phases. It demonstrated the necessity of the MCH neurons to increase REM sleep during warming toward 'just right' room temperature. Credit: Pascal Gugler for Insel Gruppe AG

Every night while sleeping, we cycle between two very different states of

sleep. Upon falling asleep, we enter non-rapid eye movement (non-REM) sleep where our breathing is slow and regular and movement of our limbs or eyes are minimal. Approximately 90 minutes later, however, we enter rapid eye movement (REM) sleep. This is a paradoxical state where our breathing becomes fast and irregular, our limbs twitch, and our eyes move rapidly. In REM sleep, our brain is highly active, but we also become paralyzed and we lose the ability to thermoregulate or maintain our constant body temperature. "This loss of thermoregulation in REM sleep is one of the most peculiar aspects of sleep, particularly since we have finely-tuned mechanisms that control our body temperature while awake or in non-REM sleep," says Markus Schmidt of the Department for BioMedical Research (DBMR) of the University of Bern, and the Department of Neurology, Inselspital, Bern University Hospital. On the one hand, the findings confirm a hypothesis proposed earlier by Schmidt, senior author of the study, and on the other hand represent a breakthrough for sleep medicine. The paper was published in *Current Biology* and highlighted by the editors with a comment.

## **A control mechanism saving energy**

The need to maintain a constant body temperature is our most expensive biological function. Panting, piloerection, sweating, or shivering are all energy consuming body reactions. In his hypothesis, Markus Schmidt suggested that REM sleep is a behavioral strategy that shifts energy resources away from costly thermoregulatory defense toward, instead, the brain to enhance many brain functions. According to this energy allocation hypothesis of sleep, mammals have evolved mechanisms to increase REM sleep when the need for defending our body temperature is minimized or, rather, to sacrifice REM sleep when we are cold. "My hypothesis predicts that we should have neural mechanisms to dynamically modulate REM sleep expression as a function of our room temperature," says Schmidt. Neuroscientists at the DBMR at the University of Bern and the Department of Neurology at Inselspital, Bern

University Hospital, have now confirmed his hypothesis and found neurons in the hypothalamus that specifically increase REM sleep when the room temperature is "just right."



Markus H. Schmidt, MD, PhD, University of Bern and Inselspital, Bern University Hospital. Credit: Pascal Gugler for Insel Gruppe AG

## **REM sleep promoting neurons**

The researchers discovered that a small population of neurons within the hypothalamus, called melanin-concentrating hormone (MCH) neurons, play a critical role in how we modulate REM sleep expression as a function of ambient (or room) temperature. The researchers showed that mice will dynamically increase REM sleep when the room temperature

is warmed to the high end of their comfort zone, similar to what has been shown for human sleep. However, genetically engineered mice lacking the receptor for MCH are no longer able to increase REM sleep during warming, as if they are blind to the warming temperature. The authors used optogenetics techniques to specifically turn on or off MCH neurons using a laser light time locked to the temperature warming periods. Their work confirms the necessity of the MCH system to increase REM sleep when the need for body temperature control is minimized.

## Breakthrough for sleep medicine

This is the first time that an area of the brain has been found to control REM sleep as a function of room [temperature](#). "Our discovery of these neurons has major implications for the control of REM sleep," says Schmidt. "It shows that the amount and timing of REM sleep are finely tuned with our immediate environment when we do not need to thermoregulate. It also confirms how dream sleep and the loss of thermoregulation are tightly integrated."

REM sleep is known to play an important role in many brain functions such as memory consolidation. REM sleep comprises approximately one quarter of our total sleep time. "These new data suggest that the function of REM sleep is to activate important brain functions specifically at times when we do not need to expend energy on thermoregulation, thus optimizing use of energy resources," says Schmidt.

**More information:** Noémie Komagata et al, Dynamic REM Sleep Modulation by Ambient Temperature and the Critical Role of the Melanin-Concentrating Hormone System, *Current Biology* (2019). [DOI: 10.1016/j.cub.2019.05.009](https://doi.org/10.1016/j.cub.2019.05.009)

Provided by University of Bern

Citation: 'Goldilocks' neurons promote REM sleep (2019, June 19) retrieved 4 May 2024 from <https://medicalxpress.com/news/2019-06-goldilocks-neurons-rem.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.