

Higher salt intake can cause gastrointestinal bloating

June 27 2019



Credit: CC0 Public Domain

A study led by researchers at the Johns Hopkins Bloomberg School of Public Health found that individuals reported more gastrointestinal bloating when they ate a diet high in salt.

The scientists re-analyzed data from a large clinical trial—the Dietary Approaches to Stop Hypertension-Sodium trial (DASH-Sodium)—conducted two decades ago, and found that high [sodium intake](#) increased bloating among trial participants. The researchers also found that the high-fiber DASH diet increased bloating among trial participants compared to a low-fiber control diet.

The study was published June 17 in the *American Journal of Gastroenterology*.

"Bloating is one of the leading gastrointestinal complaints in the U.S. and can be exacerbated in some people by a high-fiber diet; our results suggest that they might be able to reduce that bloating, without compromising on healthy fiber, by lowering their [sodium](#) intake," says study senior author Noel Mueller, Ph.D., MPH, an assistant professor in the Department of Epidemiology at the Bloomberg School.

Bloating is estimated to affect up to a third of U.S. adults overall, and more than 90 percent of those with irritable bowel syndrome. Bloating features a buildup of excess gas in the gut. The production of gas can be attributed to gas-producing gut bacteria breaking down fiber. There is also some evidence that sodium can stimulate bloating. The study by Mueller and colleagues is the first to examine sodium as a cause of bloating in the context of low- and high-fiber diets.

The study analyzed data from the DASH-Sodium trial, which was co-led by Bloomberg School researcher Lawrence Appel, MD, MPH, and sponsored by the National Heart, Lung and Blood Institute. Conducted at four clinical centers during 1998-99, it tested the DASH diet, a high-fiber diet which is relatively low in fat and high in fruits, nuts, and vegetables, against a low-fiber control diet. Each of the two diets was tested at three levels of sodium, and the 412 participants all had [high blood pressure](#) at the trial start. The trial was set up chiefly to determine

the effect of dietary sodium and other factors on blood pressure, but included data on participants' reports of bloating—data that Mueller and colleagues analyzed for the new study.

The team found that prior to the trial, 36.7 percent of the participants reported bloating, which is more or less in line with national surveys of bloating prevalence. They found too that the high-fiber DASH diet increased the risk of bloating by about 41 percent, compared to the low-fiber control diet—and men were more susceptible to this effect, compared to women. But the scientists also determined that sodium was a factor in bloating. When they combined data from the DASH and control diets, and compared the highest level of sodium intake to the lowest, they found that the high-sodium versions of those diets collectively increased the risk of bloating by about 27 percent compared to the low-sodium versions.

The key implication is that reducing sodium can be an effective way to reduce bloating—and in particular may be able to help people maintain a healthy, high-fiber [diet](#).

How sodium causes bloating is still being studied. Salt causes water retention, and that may be one factor. "We hypothesize that sodium intake also alters the [gut microbiome](#) in a manner that modifies bacterial sulfide production," Mueller says.

He and his team are now researching how bloating is affected by the major dietary macronutrients: protein, carbs and fat.

More information: Allison W. Peng et al. Effects of the DASH Diet and Sodium Intake on Bloating, *The American Journal of Gastroenterology* (2019). [DOI: 10.14309/ajg.0000000000000283](https://doi.org/10.14309/ajg.0000000000000283)

Provided by Johns Hopkins University Bloomberg School of Public Health

Citation: Higher salt intake can cause gastrointestinal bloating (2019, June 27) retrieved 6 May 2024 from <https://medicalxpress.com/news/2019-06-higher-salt-intake-gastrointestinal-bloating.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.