

More support needed to increase HIV testing in GP practices

June 10 2019



Credit: CC0 Public Domain

One-off training sessions for GPs are not enough to increase rates of HIV testing in general practice and greater support is needed, according to researchers from the University of Bristol's Centre for Academic

Primary Care and National Institute for Health Research (NIHR) Health Protection Research Unit in Evaluation of Interventions published in BMC Family Practice.

HIV-infected patients often present to their GP several times with signs, symptoms or a diagnosis more commonly found in those with HIV before being diagnosed, but healthcare professionals often miss this opportunity to test for HIV.

The researchers evaluated an education intervention designed to encourage GPs in practices with higher than average rates of HIV (more than two cases per 1,000 population) to undertake more routine HIV testing. Staff from 19 GP practices in Bristol, South Gloucestershire and North Somerset took part.

The intervention consisted of a one-hour interactive training workshop delivered by a specialist sexual health clinician. This included information on HIV testing recommendations, common signs and symptoms, who to test, how to discuss the test with patients, overcoming barriers to testing, how to manage negative and positive results, and linkage to care. The training concluded with a discussion, supported by case study examples, about ways to increase practice-level testing.

The researchers found that although there was a small increase in HIV testing in the six months after the training workshop, it could not be attributed to the intervention. Staff were positive about the intervention and reported that it increased their knowledge of HIV and skills in offering a HIV [test](#).

A possible explanation for the lack of effect on HIV testing rates is that the training was delivered as a one-off session. Some clinicians commented that they struggled to remember the training content beyond the key messages. Also, the focus of the intervention was on information

rather than skills-based [training](#) which may have failed to overcome barriers experienced by staff in offering HIV tests, such as lack of time, managing patient expectations and not prioritising HIV testing.

Organisational and structural barriers included severely limited resources available for delivering the [intervention](#) due to local authority budget restrictions.

Dr. Jo Kesten from the NIHR Health Protection Research Unit in Evaluation of Interventions at the University of Bristol and NIHR CLAHRC West, and a lead author of the study, said: "Increasing the uptake of HIV testing across healthcare settings and reducing the stigma surrounding HIV testing is a major priority. A single educational workshop on HIV testing, although experienced positively, is likely to require repetition and support from additional measures to help encourage increased HIV testing rates. Our research highlights several continued barriers to testing suggesting that more support is needed to increase clinicians' motivation and opportunity for HIV testing in primary care."

Dr. Mark Gompels, Immunology Consultant at North Bristol NHS Trust said: "It is vital that we support GPs to increase HIV testing in high-prevalence areas. Over 100,000 people in the UK are living with HIV, 96 percent of whom are on antiretroviral therapy or ART. Of those on ART, 94 percent have a consistently undetectable HIV viral load and therefore cannot transmit HIV to a sexual partner. This is fantastic news. However, around 10,400 people are undiagnosed and unaware they are HIV positive and are at greater risk of passing the virus on to others through unprotected sex. Also, individuals who are unaware they are HIV positive may only be diagnosed once their infection is advanced and they have become seriously ill. It is vital we determine the most effective methods of identifying those who unknowingly have HIV infections."

More information: Charlotte F. Davies et al. Evaluation of an educational intervention to increase HIV-testing in high HIV prevalence general practices: a pilot feasibility stepped-wedged randomised controlled trial, *BMC Family Practice* (2018). [DOI: 10.1186/s12875-018-0880-9](https://doi.org/10.1186/s12875-018-0880-9)

Joanna M. Kesten et al. Qualitative evaluation of a pilot educational intervention to increase primary care HIV-testing, *BMC Family Practice* (2019). [DOI: 10.1186/s12875-019-0962-3](https://doi.org/10.1186/s12875-019-0962-3)

Provided by University of Bristol

Citation: More support needed to increase HIV testing in GP practices (2019, June 10) retrieved 17 May 2024 from <https://medicalxpress.com/news/2019-06-hiv-gp.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|