

## **Improvements in insulin release wane after treatment stops in adults with type 2 diabetes**

June 10 2019



Dr. Steven Kahn, study chair for the Restoring Insulin Secretion (RISE) Consortium, talks with RISE participant Faamafi Faamafi Jr. about his experience in the adult medication trial. Kahn is an endocrinologist at VA Puget Sound Health Care System and University of Washington School of Medicine, both in Seattle. The site was one of eight RISE locations across the country looking at type 2 diabetes treatments in children and adults. Credit: Christopher Pacheco, VA Puget Sound Health Care System



A set of clinical trials examining youth and adults with type 2 diabetes or impaired glucose tolerance has found that disease progression in adults slowed during medical treatment but resumed after treatment stopped. Youth on the same treatment had markedly poorer outcomes with continued disease progression both during and after the treatment. This research, funded by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), was published June 9 in the journals *Diabetes* and *Diabetes Care* and presented at the American Diabetes Association Scientific Sessions in San Francisco. NIDDK is part of the National Institutes of Health.

The Restoring Insulin Secretion (RISE) Adult and Pediatric Medication Studies compared the use of different treatments among adults aged 20-65 and youth aged 10-19 with impaired glucose tolerance or early onset type 2 diabetes with the aim of preserving beta cell function—key to the body's ability to make and release insulin.

In the adult study, participants were randomly assigned to receive either long-acting insulin (glargine) for three <u>months</u> followed by nine months of metformin, the drug liraglutide in combination with metformin for 12 months, metformin alone for 12 months, or a placebo. Researchers wanted to see whether early aggressive treatment could have a lasting effect on slowing or stopping the decline of beta cell function that occurs in people with type 2 diabetes. Participants were monitored for three additional months after treatments ended.

Results showed that adult participants had improvements in beta cell function and blood glucose control while on the treatments, with those in the liraglutide plus metformin group showing the most improvement after 12 months. However, these improvements did not persist among any of the groups after treatment ended.



"The RISE medication studies show that these treatments for type 2 diabetes do not make lasting changes to beta cells," said study chair Dr. Steven Kahn from the Veteran's Affairs Puget Sound Health Care System and University of Washington School of Medicine. "For adults, we saw that the treatment options were equally effective while people were actively on them—but people need to stay on treatment to maintain the benefits."

In a companion paper, findings from the adult study were compared to those from the RISE Pediatric Medication Study <u>published in 2018</u>, which showed that beta cell function declined in the two youth treatment groups during active treatment, and worsened after treatment ended.

The RISE Adult and Pediatric Medication Studies were designed together to enable direct comparison of the effect of treatment on youth and adults. The youth study compared the use of three months of insulin glargine followed by metformin for nine months to metformin alone for 12 months. Insulin glargine and metformin are the only U.S. Food and Drug Administration-approved medications for youth with type 2 diabetes.

"Though the medications' effectiveness for adults while on treatment is reassuring, the poor results for youth in the study, both during and after treatment, underscore the continued urgent need for new approaches to prevent and treat type 2 diabetes in youth, since the disease progresses more rapidly given the same treatment as adults," said Dr. Ellen Leschek, a study author and the NIDDK project scientist for RISE.

This close comparison between adults and youth receiving the same type 2 diabetes treatments supports earlier research suggesting the disease is more aggressive in youth than adults, and points to new areas of research that may help explain why.



"The RISE studies show that type 2 diabetes affects youth differently, and more aggressively, than <u>adults</u>, said NIDDK Director Dr. Griffin P. Rodgers. "These findings demonstrate the need for continued research to identify new treatment strategies to control and treat type 2 <u>diabetes</u>, and underscore the need to focus on prevention efforts, especially for <u>youth</u>."

**More information:** undefined undefined. Effects of Treatment of Impaired Glucose Tolerance or Recently Diagnosed Type 2 Diabetes With Metformin Alone or in Combination With Insulin Glargine on  $\beta$ -Cell Function: Comparison of Responses In Youth And Adults, *Diabetes* (2019). DOI: 10.2337/db19-0299

undefined undefined. Lack of Durable Improvements in β-Cell Function Following Withdrawal of Pharmacological Interventions in Adults With Impaired Glucose Tolerance or Recently Diagnosed Type 2 Diabetes, *Diabetes Care* (2019). DOI: 10.2337/dc19-0556

Provided by National Institute of Diabetes and Digestive and Kidney Diseases

Citation: Improvements in insulin release wane after treatment stops in adults with type 2 diabetes (2019, June 10) retrieved 2 May 2024 from https://medicalxpress.com/news/2019-06-insulin-wane-treatment-adults-diabetes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.