

# Study finds micronutrient deficiencies common at time of celiac disease diagnosis

June 24 2019

---



Credit: Ragesoss/Wikipedia

Micronutrient deficiencies, including vitamins B12 and D, as well as folate, iron, zinc and copper, are common in adults at the time of diagnosis with celiac disease. These deficiencies should be addressed at that time, according to a study by Mayo Clinic researchers.

The retrospective study of 309 adults newly diagnosed with [celiac disease](#) at Mayo Clinic from 2000 to 2014 also found that low body weight and [weight loss](#), which are commonly associated with celiac disease, were less common. Weight loss was seen in only 25.2% of patients, and the average body mass index was categorized as overweight. The study will appear in the July issue of *Mayo Clinic Proceedings*.

"It was somewhat surprising to see the frequency of micronutrient deficiencies in this group of newly diagnosed patients, given that they were presenting fewer symptoms of malabsorption," says Adam Bledsoe, M.D., a gastroenterology fellow at Mayo Clinic's Rochester campus.

Celiac disease is an immune reaction to consuming gluten, a protein found in wheat, barley and rye. Eating gluten triggers an immune response in the [small intestine](#) that over time damages the intestine's lining and prevents it from absorbing some nutrients, leading to diarrhea, fatigue, anemia, weight loss and other complications.

Based on recent data, the prevalence of celiac disease in the U.S. is 1 in 141 people, and its prevalence has increased over the past 50 years.

"Our study suggests that the presentation of celiac disease has changed from the classic weight loss, anemia and diarrhea, with increasing numbers of patients diagnosed with nonclassical symptoms," says Dr. Bledsoe, the study's primary author. "Micronutrient deficiencies remain common in adults, however, and should be assessed." Assessment should include vitamin D, iron, folic acid, vitamin B12, zinc and copper.

Zinc deficiency was observed most frequently at diagnosis, the study says, with 59.4% of patients having a deficiency. Other deficiencies included iron, vitamin D, copper, vitamin B12 and folate.

The nutritional deficiencies have potential health ramifications, though in this [retrospective study](#) the clinical implications remain unknown. "Further studies are needed to better define the implications of the deficiencies, optimal replacement strategies and follow-up," says Dr. Bledsoe.

Provided by Mayo Clinic

Citation: Study finds micronutrient deficiencies common at time of celiac disease diagnosis (2019, June 24) retrieved 27 April 2024 from <https://medicalxpress.com/news/2019-06-micronutrient-deficiencies-common-celiac-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.