

Opioid analgesics increase the risk of pneumonia among persons with Alzheimer's disease

June 4 2019



Credit: CC0 Public Domain

Opioid analgesics were associated with a 30 percent increase in the risk of pneumonia in persons with Alzheimer's disease, a recent study from the University of Eastern Finland shows. The risk was most pronounced

in the first two months of use. This is the first study to investigate the association between opioids and pneumonia in this population. The results were published in the *Journal of Alzheimer's Disease*.

The risk of pneumonia was highest among those using [strong opioids](#), such as oxycodone or fentanyl, but the risk was also increased among those using buprenorphine, tramadol or codeine. Persons with active cancer were excluded from the study, and other diseases and drug use were accounted for in the analyses. Opioids impair the cough reflex and respiratory functions and cause sedation, possibly underlying the increased risk for pneumonia.

Pneumonia is a severe but relatively common complication among persons with Alzheimer's disease, and often leads to hospitalization and even death. Opioids are important in the treatment of severe acute pain, but treatment should be initiated at a low dose and regularly assessed for both benefit and harm. The diagnosis and assessment of pain is challenging among persons with Alzheimer's disease due to problems in communication.

The study is part of the MEDALZ cohort, which included 5,623 persons with Alzheimer's disease diagnosed in Finland during 2010-2011. Each person initiating opioid use was matched with a comparison person with Alzheimer's disease who did not initiate opioid use but had the same age, gender and year of Alzheimer's diagnosis. Data for the study were derived from Finnish nationwide registers.

More information: Aleksi Hamina et al. Hospital-Treated Pneumonia Associated with Opioid Use Among Community Dwellers with Alzheimer's Disease, *Journal of Alzheimer's Disease* (2019). [DOI: 10.3233/JAD-181295](https://doi.org/10.3233/JAD-181295)

Provided by University of Eastern Finland

Citation: Opioid analgesics increase the risk of pneumonia among persons with Alzheimer's disease (2019, June 4) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2019-06-opioid-analgesics-pneumonia-persons-alzheimer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.