

Poor sleep is unlikely to contribute to higher BMI in children

June 5 2019



Credit: CC0 Public Domain

Children need more than a good night's sleep to have a healthy weight,



according to a new study in the journal *Obesity* that explores the relationship between sleep, body mass index (BMI) and cortisol levels in children.

Shorter sleep duration has been linked to higher BMI, and that relationship has led researchers to consider whether poor sleep might contribute to higher BMI. This study found that kids with shorter sleep duration had higher BMI in general, but no evidence of a causal link.

"Surprisingly, our findings actually suggest that changes in sleep at one age did not lead to changes in BMI later in childhood, or vice versa," said Kristine Marceau, an assistant professor of human development and <u>family studies</u> at Purdue University. "Instead, the relationship between BMI and sleep was stable over time and was likely caused by other factors in kids' lives, such as genetics, family routines and habits. This means that we would expect that interventions that are aimed only at improving sleep habits in <u>children</u> may not be as effective for changes in BMI as interventions that can target the factors that lead to changes in both sleep and BMI. More research is needed to identify those targets."

The researchers also looked at <u>cortisol</u>, a stress hormone that works with certain parts of the brain to help regulate a number of bodily functions, including metabolism and blood sugar levels, which could explain the link between sleep length and higher BMI. There was no link with cortisol levels at a younger age, but the study did indicate an emerging relationship between cortisol and BMI during middle childhood, which was independent of sleep.





Kristine Marceau, assistant professor of human development and family studies. Credit: Purdue University

"We did not find that <u>cortisol levels</u> were very stable across childhood," Marceau said. "Cortisol production is highly variable during <u>early</u> <u>childhood</u>, suggesting that cortisol patterns may still be developing at this age."

According to Marceau, most studies looking at stress hormone levels and BMI involve adults. Marceau, who specializes in gene-environment interplay, physiological responses to stress, and pubertal development, and the research team looked at data on 361 children ages 4 ½ to 9 years old, which originated from a long-term study of adopted children who are now being assessed in adolescence. This is the first study of its type, and Marceau points out that the study needs to be replicated.



Parents kept children's sleeping logs, which showed the children were averaging about 10 hours of sleep a night. The BMI range fell within the those recommended by the Centers for Disease Control and Prevention.

More information: Kristine Marceau et al. Longitudinal Associations of Sleep Duration, Morning and Evening Cortisol, and BMI During Childhood, *Obesity* (2019). DOI: 10.1002/oby.22420

Provided by Purdue University

Citation: Poor sleep is unlikely to contribute to higher BMI in children (2019, June 5) retrieved 27 April 2024 from https://medicalxpress.com/news/2019-06-poor-contribute-higher-bmi-children.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.