

Repurposing existing drugs or combining therapies could help in the treatment of autoimmune diseases

June 17 2019



Credit: CC0 Public Domain

Research led by the University of Birmingham has found re-purposing already existing drugs or combining therapies could be used to treat patients who have difficult to treat autoimmune diseases.

Funded by Versus Arthritis, the research was led by the University of Birmingham's Institute of Inflammation and Ageing and Institute of Cardiovascular Sciences and was published today (June 17th) in *Proceedings of the National Academy of Sciences*.

The research, a collaboration with the University of Oxford, University of Cambridge, University of York, Université Rennes in France, and the University of Lausanne in Switzerland, was supported by the National Institute for Health Research Birmingham Biomedical Research Centre.

Dr. Saba Nayar, of the University of Birmingham, explained: "In this study, we found for the first time that fibroblasts—[cells](#) that play a critical role in healing—also play a key role in the process of the formation of tertiary lymphoid structures, which are small clusters of blood and tissue cells found at the sites of chronic inflammation.

"Inflammation is the body's process of fighting against things that harm it, such as infections, injuries, and toxins, in an attempt to heal itself. When something damages cells, our bodies releases chemicals that trigger a response from our immune system.

"This response usually lasts for a few hours or days in the case of acute inflammation, however in [chronic inflammation](#) the response lingers, leaving your body in a constant state of alert. Chronic [inflammation](#) occurs in a range of conditions from cancer to arthritis and autoimmune conditions—illnesses or disorders that occur when [healthy cells](#) get destroyed by the body's own immune system."

Dr. Joana Campos, also of the University of Birmingham, added: "Tertiary lymphoid structures are believed to play a key role in the progression of autoimmune conditions such as Sjögren's Syndrome—a condition that affects parts of the body that produce fluids like tears and spit.

"Previously research has not identified the role fibroblasts play in the formation and maintenance of tertiary lymphoid structures.

"We proved that fibroblasts expand and acquire immunological features in a process that is dependent on two cytokines—substances which are secreted by cells including fibroblasts in the [immune system](#)."

Dr. Francesca Barone, also of the University of Birmingham, said: "Our research has led us to conclude that, by re-purposing already existing drugs or combining therapies, we could use these medications to directly target immune cells and fibroblasts to attack these cytokines in patients who have difficult to treat [autoimmune diseases](#) in which the formation of tertiary lymphoid structures plays a critical role.

"Our findings were surprising and unexpected and have addressed functional questions that the science community has been trying to address since tertiary lymphoid structures were first discovered."

More information: Saba Nayar et al., "Immunofibroblasts are pivotal drivers of tertiary lymphoid structure formation and local pathology," *PNAS* (2019). www.pnas.org/cgi/doi/10.1073/pnas.1905301116

Provided by University of Birmingham

Citation: Repurposing existing drugs or combining therapies could help in the treatment of autoimmune diseases (2019, June 17) retrieved 30 April 2024 from <https://medicalxpress.com/news/2019-06-repurposing-drugs-combining-therapies-treatment.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--