

Signature of an ineffective immune response to cancer revealed

June 20 2019



Credit: CC0 Public Domain

Our immune system is programmed to destroy cancer cells. Sometimes it has trouble slowing disease progression because it doesn't act quickly or strongly enough. In a study published in *The Journal of Clinical Investigation*, researchers from the University of Montreal Hospital

Research Centre (CRCHUM) revealed the genetic signature of this failed immune response for the first time.

"This distinctive signature includes 28 genes. We were able to identify the signature by studying the genetic programming of immune cells from kidney tumours and from the blood of patients whose cancer was progressing, and then comparing it with the genetic programming of healthy people's immune cells. This signature could help us predict which patients will fare worse," said Réjean Lapointe, researcher and head of the cancer research theme at the CRCHUM.

The research team also showed that these same genes that are linked to a failed [immune response](#) are found in patients with cancers other than kidney cancer or with bacterial and viral infections such as AIDS.

"Thanks to our sample bank and powerful bioinformatics tools, we were able to validate the relevance of this signature on nearly 11,500 patients. Those with the worst clinical outcome had a higher presence of these 28 genes in their immune cells," explained Lapointe, who is also a professor at the Université de Montréal and the scientific director of the Institut de cancer de Montréal.

Of the 28 genes identified among the 20,000 coding genes in the [human body](#), the scientists were able to show that the expression of three [specific genes](#) may even be able to predict a person's chance of survival.

A very detailed mapping

To achieve such conclusive results, Lapointe's team had to map the differences in genetic programming between the immune cells found in tumours and those circulating in the bloodstream, as well as the differences between [immune cells](#) that had infiltrated the tumours of patients with a very aggressive illness and those who live longer.

"There are already a few articles in the scientific literature that partly replicate some of our results. A very recent HIV study shows that the MMP9 protein is associated with patients who do not control the infection very well. That's what we had concluded in our study," said Lapointe.

This massive basic research undertaking will help identify the therapeutic targets or mechanisms that can be "unlocked" in order to wake up the [immune system](#) and eliminate [cancer cells](#).

More information: Anne Monette et al, Failed immune responses across multiple pathologies share pan-tumor and circulating lymphocytic targets, *Journal of Clinical Investigation* (2019). [DOI: 10.1172/JCI125301](#)

Provided by University of Montreal

Citation: Signature of an ineffective immune response to cancer revealed (2019, June 20) retrieved 25 June 2024 from <https://medicalxpress.com/news/2019-06-signature-ineffective-immune-response-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--