

Why do women live longer than men?

June 10 2019, by Melinda Martin-Khan



Credit: AI-generated image (disclaimer)

In Australia, an average baby boy born in 2016 could expect to live to 80, while a baby girl born at the same time could expect to live until closer to 85. A similar gap in life expectancy between men and women is seen around the world.

As we better understand why people die, we're learning how biological and behavioral factors may partly explain why <u>women</u> live longer than



men.

Scientific advancements also impact the health of women and men differently.

Biology and behavior

While women may live longer than men, they report more illnesses, more doctor visits and more hospital stays than men. This is known as the <u>morbidity-mortality paradox</u> (that is, women are sicker but live longer).

One explanation is that women suffer from <u>illnesses less likely to kill</u> them. Examples of chronic non-fatal illnesses more common in women include <u>migraines</u>, <u>arthritis</u> and <u>asthma</u>. These conditions may lead to poorer health, but don't increase a woman's risk of premature or early death.

But men are more susceptible to health conditions that can kill them. For example, men tend to have more fat surrounding their organs (called visceral fat) and women tend to have more fat under their skin (called subcutaneous fat). Visceral fat is a risk factor for coronary heart disease, the leading underlying cause of death for Australian men.

Coronary heart disease, which results from a combination of biological factors and lifestyle habits, is a major reason for the difference in mortality between men and women.

Other biological factors may contribute to men aging faster than women, but these remain to be fully understood. For example, <u>testosterone in men</u> contributes to their generally larger bodies and deeper voices. In turn, this <u>may accelerate the age-related changes</u> in their bodies compared to women.



On the flip side, women may have a slight advantage from <u>protective</u> <u>factors connected with estrogen</u>. Coronary heart disease has been observed as three times lower in women than in men before menopause, but not after, indicating that endogenous oestrogens could have a protective effect in women.

Some behaviors that can lead to an earlier death are more common in men. Accidental deaths, including those caused by assault, poisoning, transport accidents, falls and drownings, are particularly high among young males aged 15-24.

Leading causes of death

Leading underlying causes of death, by sex, Australia (2016)

	Male	Female
Coronary heart disease	10,870	8,207
Dementia and Alzheimer's disease	4,679	8,447
Cerebrovascular disease (stroke)	4,239	6,212
Lung cancer	5,023	3,387
Chronic obstructive pulmonary disorde (lung disease)	3,903	3,309

Credit: The Conversation

Men also have <u>a greater tendency</u> to smoke, eat poorly and avoid exercise. These habits lead to <u>often fatal chronic illnesses</u>, including <u>stroke</u> and <u>type 2 diabetes</u>, and are also <u>risk factors for dementia</u>.

Developments in science and public health



Many <u>scientific discoveries</u> have led to improved <u>clinical practice</u> or changes in <u>government health policies</u> that have benefited the <u>lives of women</u>.

For example, <u>innovations in birth control</u> have enabled greater choice and control over family size and timing. This has resulted in fewer pregnancies that may have led to dangerous births, and improved general physical and <u>mental health for women</u>. Improved clinical care has resulted in fewer women dying during childbirth.

Public health programs such as <u>screening for breast cancer</u> have had impacts on life expectancy over time. Similarly, <u>vaccines to prevent</u> <u>cervical cancer</u> have now been distributed in 130 countries.

Of course, there have been similar <u>public health</u> policies and clinical innovations that have benefited men too, <u>like screening for bowel caner</u>.

So although we may have some insights, we can't conclusively answer why women continue to live longer than men.

Mind the gap

The gap between men and women <u>decreases the longer they live</u>. In 2016, at birth in Australia, the gap was 4.2 years, with a male expected to die at 80 on average. But as that male gets older, the gap decreases to 2.7 years at age 65, to one year at age 85 and to just 0.3 years at age 95.

This suggests men who live to an older age have been able to avoid certain health risks, giving them a greater prospect of a longer life.

Ultimately, none of us have control of when or how we're going to die. But paying attention to factors that we can change (such as maintaining a healthy diet, doing exercise and avoiding smoking) can <u>reduce the risk</u> of



dying earlier from a preventable chronic disease.

While women may always live longer than men, by a year or two, men can try to make some lifestyle changes to reduce this gap. That being said, women should work towards these goals for a long and healthy life, too.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Why do women live longer than men? (2019, June 10) retrieved 26 June 2024 from https://medicalxpress.com/news/2019-06-women-longer-men.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.