


Zebrafish capture a 'window' on the cancer process

June 4 2019



An immune cell (magenta) breaches the basement membrane to access cancer cells. The basement membrane separates different tissues in the body and plays an important role as a barrier in confining tumours. Credit: University of Bristol

Cancer-related inflammation impacts significantly on cancer


development and progression. New research has observed in zebrafish, for the first time, that inflammatory cells use weak spots or micro-perforations in the extracellular matrix barrier layer to access skin cancer cells.

The research, led by the University of Bristol and published in *Cell Reports* Tuesday 4 June, used translucent zebrafish to model several sorts of skin cancer and live image how inflammatory cells find the growing cancer cells in the skin. To access the cancer cells, the [immune cells](#) need to first breach an extracellular matrix barrier layer called the basement membrane zone.

The researchers observed weak spots in the basement membrane zone, which the inflammatory cells use as easy routes to access the cancer. Those clones of cancer cells nearest to the weak spots tend to receive more inflammatory cell visits and as a consequence they grow faster.

Paul Martin, Professor of Cell Biology in the Schools of Biochemistry and Physiology, Pharmacology & Neuroscience at the University of Bristol, said: "As the zebrafish are translucent, we can watch inflammatory cells interacting with cancer cells in ways not possible in our own tissues.

"This 'window' on the cancer process has revealed 'weak spots' in the barrier layer that inflammatory cells must breach in order to access and feed the cancer cells within the skin. Now we know these micro-perforations exist we can target them with cancer therapeutics."



Micrograph of a region of zebrafish skin where a track of cancer cells has disrupted the epithelium much like a mole burrowing beneath a lawn of grass. Credit: University of Bristol

Inflammatory cells ([white blood cells](#)) are known to be key players encouraging cancer malignancy from the earliest stages of cancer initiation, but it has been unclear how they gain access to the [cancer cells](#) which generally develop within epithelial tissues, which have a barrier layer of extracellular matrix between them and the tissues from where the inflammatory cells arise. Inflammatory cells could kill the growing cancer but instead tend to nourish it and encourage cancer progression.

The study's findings have clear clinical relevance to cancer patients because microperforations in the basement membrane zone have been shown to occur in human airways and guts and so could act as similar portals to let [inflammatory cells](#) gain access to the cancer.

More information: Proteolytic and Opportunistic Breaching of the Basement Membrane Zone by Immune Cells during Tumor Initiation, *Cell Reports* (2019). [DOI: 10.1016/j.celrep.2019.05.029](https://doi.org/10.1016/j.celrep.2019.05.029) , [www.cell.com/cell-reports/full ... 2211-1247\(19\)30640-0](https://www.cell.com/cell-reports/full...2211-1247(19)30640-0)

Provided by University of Bristol

Citation: Zebrafish capture a 'window' on the cancer process (2019, June 4) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-06-zebrafish-capture-window-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.